

Exhibit 29

**IN THE UNITED STATES DISTRICT COURT
FOR THE SOUTHERN DISTRICT OF WEST VIRGINIA
CHARLESTON DIVISION**

B.P.J., by her next friend and mother,
HEATHER JACKSON,

Plaintiff,

vs.

WEST VIRGINIA STATE BOARD OF
EDUCATION; HARRISON COUNTY BOARD
OF EDUCATION; WEST VIRGINIA
SECONDARY SCHOOLS ACTIVITIES
COMMISSION; W. CLAYTON BURCH, in his
official capacity as State Superintendent, DORA
STUTLER, in her official capacity as the
Harrison County Superintendent, and the
STATE OF WEST VIRGINIA,

Defendants,

and

LAINIEY ARMISTEAD,

Defendant-Intervenor.

Case No. 2:21-cv-00316

Hon. Joseph R. Goodwin

DECLARATION OF GREGORY A. BROWN, PH.D., FACSM

I, Dr. Gregory A. Brown, pursuant to 28 U.S. Code § 1746, declare under penalty of perjury under the laws of the United States of America that the facts contained in my Expert Declaration of Gregory A. Brown, Ph.D., FACSM in the Case of B.P.J. v. West Virginia State Board of Education, attached hereto, are true and correct to the best of my knowledge and belief, and that the opinions expressed therein represent my own expert opinions.

Executed on February 23, 2022.



Gregory A. Brown

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Expert Report of
Gregory A Brown, Ph.D. FACSM
In the case of B.P.J. vs. West Virginia State Board of Education.

Table of Contents

Table of Contents.....	iv
Personal Qualifications and Disclosure.....	1
Overview	4
I. The scientific reality of biological sex.....	5
II. Biological men, or adolescent boys, have large, well-documented performance advantages over women and adolescent girls in almost all athletic contests.....	7
A. Men are stronger.	10
B. Men run faster.....	12
C. Men jump higher and farther.	15
D. Men throw, hit, and kick faster and farther.	16
E. Males exhibit faster reaction times.	17
III. Men have large measured physiological differences compared to women which demonstrably or likely explain their performance advantages.	17
A. Men are taller and heavier than women.....	18
B. Males have larger and longer bones, stronger bones, and different bone configuration.....	18
C. Males have much larger muscle mass.....	20
D. Females have a larger proportion of body fat.	20
E. Males are able to metabolize and release energy to muscles at a higher rate due to larger heart and lung size, and higher hemoglobin concentrations.	21
IV. The role of testosterone in the development of male advantages in athletic performance	23
A. Boys exhibit advantages in athletic performance even before puberty. ..	25

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

B.	The rapid increase in testosterone across male puberty drives characteristic male physiological changes and the increasing performance advantages.	37
V.	The available evidence shows that suppression of testosterone in a male after puberty has occurred does not substantially eliminate the male athletic advantage.....	39
A.	Empirical studies find that males retain a strong performance advantage even after lengthy testosterone suppression.....	39
B.	Testosterone suppression does not reverse important male physiological advantages.	46
C.	Responsible voices internationally are increasingly recognizing that suppression of testosterone in a male after puberty has occurred does not substantially reverse the male athletic advantage.	50
	Conclusions	56
	Bibliography	a
	Appendix 1 – Data Tables	h
	Presidential Physical Fitness Results.....	h
	Data Compiled from Athletic.Net.....	j
	Appendix 2 – Scholarly Publications in Past 10 Years	n

Personal Qualifications and Disclosure

I serve as Professor of Exercise Science in the Department of Kinesiology and Sport Sciences at the University of Nebraska Kearney, where I teach classes in Exercise Physiology among other topics. I am also the Director of the General Studies program. I have served as a tenured (and nontenured) professor at universities since 2002.

In August 2002, I received a Doctor of Philosophy degree from Iowa State University, where I majored in Health and Human Performance, with an emphasis in the Biological Bases of Physical Activity. In May 1999, I received a Master of Science degree from Iowa State University, where I majored in Exercise and Sport Science, with an emphasis in Exercise Physiology.

I have received many awards over the years, including the Mortar Board Faculty Excellence Honors Award, College of Education Outstanding Scholarship / Research Award, and the College of Education Award for Faculty Mentoring of Undergraduate Student Research. I have authored more than 40 refereed publications and more than 50 refereed presentations in the field of Exercise Science. I have authored chapters for multiple books in the field of Exercise Science. And I have served as a peer reviewer for over 25 professional journals, including *The American Journal of Physiology*, the *International Journal of Exercise Science*, the *Journal of Strength and Conditioning Research*, and *The Journal of Applied Physiology*.

My areas of research have included the endocrine response to testosterone prohormone supplements in men and women, the effects of testosterone prohormone supplements on health and the adaptations to strength training in men, the effects of energy drinks on the physiological response to exercise, and assessment of various athletic training modes in males and females. Articles that I have published that are closely related to topics that I discuss in this white paper include:

- Studies of the effect of ingestion of a testosterone precursor on circulating testosterone levels in young men. Douglas S. King, Rick L. Sharp, Matthew D. Vukovich, Gregory A. Brown, et al., *Effect of Oral Androstenedione on Serum Testosterone and Adaptations to Resistance Training in Young Men: A Randomized Controlled Trial*, JAMA 281: 2020-2028 (1999); G. A. Brown, M. A. Vukovich, et al., *Effects of Anabolic Precursors on Serum Testosterone Concentrations and Adaptations to Resistance Training in Young Men*, INT J SPORT NUTR EXERC METAB 10: 340-359 (2000).
- A study of the effect of ingestion of that same testosterone precursor on circulating testosterone levels in young women. G. A. Brown, J. C. Dewey, et

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

al., *Changes in Serum Testosterone and Estradiol Concentrations Following Acute Androstenedione Ingestion in Young Women*, HORM METAB RES 36: 62-66 (2004.)

- A study finding (among other things) that body height, body mass, vertical jump height, maximal oxygen consumption, and leg press maximal strength were higher in a group of physically active men than comparably active women, while the women had higher percent body fat. G. A. Brown, Michael W. Ray, et al., *Oxygen Consumption, Heart Rate, and Blood Lactate Responses to an Acute Bout of Plyometric Depth Jumps in College-Aged Men And Women*, J. STRENGTH COND RES 24: 2475-2482 (2010).
- A study finding (among other things) that height, body mass, and maximal oxygen consumption were higher in a group of male NCAA Division 2 distance runners, while women NCAA Division 2 distance runners had higher percent body fat. Furthermore, these male athletes had a faster mean competitive running speed (~3.44 min/km) than women (~3.88 min/km), even though the men ran 10 km while the women ran 6 km. Katherine Semin, Alvah C. Stahlnecker, Kate A. Heelan, G. A. Brown, et al, *Discrepancy Between Training, Competition and Laboratory Measures of Maximum Heart Rate in NCAA Division 2 Distance Runners*, JOURNAL OF SPORTS SCIENCE AND MEDICINE 7: 455-460 (2008).
- A presentation at the 2021 American Physiological Society New Trends in Sex and Gender Medicine Conference entitled “Transwomen Competing in Women’s Sports: What We Know and What We Don’t”. I have also authored an August 2021 entry for the American Physiological Society Physiology Educators Community of Practice Blog (PECOP Blog) titled “The Olympics, Sex, and Gender in the Physiology Classroom.”

A list of my published scholarly work for the past 10 years appears as an Appendix.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Purpose of this Declaration

I have been asked by counsel for Defendant State of West Virginia and Intervenor Defendant Lainey Armistead in the matter of *B.P.J. by her next friend and mother Heather Jackson, v. State of West Virginia State Board of Education, et al.* to offer my opinions about the following: (a) whether males have inherent advantages in athletic performance over females, and if so the scale and physiological basis of those advantages, to the extent currently understood by science and (b) whether the sex-based performance advantage enjoyed by males is eliminated if feminizing hormones are administered to male athletes who identify as transgender (and in the case of prepubertal children, whether puberty blockers eliminate the advantage). In this declaration, when I use the terms “boy” or “male,” I am referring to biological males based on the individual’s reproductive biology and genetics as determined at birth. Similarly, when I use the terms “girl” or “female,” I am referring to biological females based on the individual’s reproductive biology and genetics as determined at birth. When I use the term transgender, I am referring to persons who are males or females, but who identify as a member of the opposite sex.

I have previously provided expert information in cases similar to this one in the form of a written declaration and a deposition in the case of *Soule vs. CIAC* in the state of Connecticut, and in the form of a written declaration in the case of *Hecox vs. Little* in the state of Idaho. I have not previously testified as an expert in any trials.

The opinions I express in this declaration are my own, and do not necessarily reflect the opinions of my employer, the University of Nebraska.

I have been compensated for my time serving as an expert in this case at the rate of \$150 per hour. My compensation does not depend on the outcome in the case.

Overview

In this declaration, I explore three important questions relevant to current discussions and policy decisions concerning inclusion of transgender individuals in women's athletic competitions. Based on my professional familiarity with exercise physiology and my review of the currently available science, including that contained in the many academic sources I cite in this report, I set out and explain three basic conclusions:

- At the level of (a) elite, (b) collegiate, (c) scholastic, and (d) recreational competition, men, adolescent boys, or male children, have an advantage over equally aged, gifted, and trained women, adolescent girls, or female children in almost all athletic events;
- Biological male physiology is the basis for the performance advantage that men, adolescent boys, or male children have over women, adolescent girls, or female children in almost all athletic events; and
- The administration of androgen inhibitors and cross-sex hormones to men or adolescent boys after the onset of male puberty does not eliminate the performance advantage that men and adolescent boys have over women and adolescent girls in almost all athletic events. Likewise, there is no published scientific evidence that the administration of puberty blockers to males before puberty eliminates the pre-existing athletic advantage that prepubertal males have over prepubertal females in almost all athletic events.

In short summary, men, adolescent boys, and prepubertal male children perform better in almost all sports than women, adolescent girls, and prepubertal female children because of their inherent physiological advantages. In general, men, adolescent boys, and prepubertal male children, can run faster, output more muscular power, jump higher, and possess greater muscular endurance than women, adolescent girls, and prepubertal female children. These advantages become greater during and after male puberty, but they exist before puberty.

Further, while after the onset of puberty males are on average taller and heavier than females, a male performance advantage over females has been measured in weightlifting competitions even between males and females matched for body mass.

Male advantages in measurements of body composition, tests of physical fitness, and athletic performance have also been shown in children before puberty. These advantages are magnified during puberty, triggered in large part by the higher testosterone concentrations in men, and adolescent boys, after the onset of

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

male puberty. Under the influence of these higher testosterone levels, adolescent boys and young men develop even more muscle mass, greater muscle strength, less body fat, higher bone mineral density, greater bone strength, higher hemoglobin concentrations, larger hearts and larger coronary blood vessels, and larger overall statures than women. In addition, maximal oxygen consumption ($VO_2\text{max}$), which correlates to ~30-40% of success in endurance sports, is higher in both elite and average men and boys than in comparable women and girls when measured in regard to absolute volume of oxygen consumed and when measured relative to body mass.

Although androgen deprivation (that is, testosterone suppression) may modestly decrease some physiological advantages that men and adolescent boys have over women and adolescent girls, it cannot fully or even largely eliminate those physiological advantages once an individual has passed through male puberty.

Evidence and Conclusions

I. The scientific reality of biological sex

1. The scientific starting point for the issues addressed in this report is the biological fact of dimorphic sex in the human species. It is now well recognized that dimorphic sex is so fundamental to human development that, as stated in a recent position paper issued by the Endocrine Society, it “must be considered in the design and analysis of human and animal research. . . . Sex is dichotomous, with sex determination in the fertilized zygote stemming from unequal expression of sex chromosomal genes.” (Bhargava et al. 2021 at 220). As stated by Sax (2002 at 177), “More than 99.98% of humans are either male or female.” All humans who do not suffer from some genetic or developmental disorder are unambiguously male or female.

2. Although sex and gender are used interchangeably in common conversation, government documents, and in the scientific literature, the American Psychological Association defines sex as “physical and biological traits” that “distinguish between males and females” whereas gender “implies the psychological, behavioral, social, and cultural aspects of being male or female (i.e., masculinity or femininity)” (<https://dictionary.apa.org>, accessed January 14, 2022). The concept that sex is an important biological factor determined at conception is a well-established scientific fact that is supported by statements from a number of respected organizations including, but not limited to, the Endocrine Society (Bhargava et al. 2021 at 220), the American Physiological Society (Shah 2014), the Institute of Medicine, and the National Institutes of Health (Miller 2014 at H781-82). Collectively, these and other organizations have stated that every cell has a sex

and every system in the body is influenced by sex. Indeed, “sex often influences gender, but gender cannot influence sex.” (Bhargava 2021 at 228.)

3. To further explain: “The classical biological definition of the **2 sexes** is that females have ovaries and make larger female gametes (eggs), whereas males have testes and make smaller male gametes (sperm) ... the definition can be extended to the ovaries and testes, and in this way the categories—female and male—can be applied also to individuals who have gonads but do not make gametes ... sex is dichotomous because of the different roles of each sex in reproduction.” (Bhargava 2021 at 221.) Furthermore, “sex determination begins with the inheritance of XX or XY chromosomes” (Bhargava 2021 at 221.) And, “Phenotypic sex differences develop in XX and XY embryos as soon as transcription begins. The categories of X and Y genes that are unequally represented or expressed in male and female mammalian zygotes ... cause phenotypic sex differences” (Bhargava 2021 at 222.)

4. Although disorders of sexual development (DSDs) are sometimes confused with discussions of transgender individuals, the two are different phenomena. DSDs are disorders of physical development. Many DSDs are “associated with genetic mutations that are now well known to endocrinologists and geneticists.” (Bhargava 2021 at 225) By contrast, a sense of transgender identity is usually not associated with any physical disorder, and “a clear biological causative underpinning of gender identity remains to be demonstrated.” (Bhargava 2021 at 226.)

5. Further demonstrating the biological importance of sex, Gershoni and Pietrokovski (2017) detail the results of an evaluation of “18,670 out of 19,644 informative protein-coding genes in men versus women” and reported that “there are over 6500 protein-coding genes with significant S[ex]D[ifferential] E[xpression] in at least one tissue. Most of these genes have SDE in just one tissue, but about 650 have SDE in two or more tissues, 31 have SDE in more than five tissues, and 22 have SDE in nine or more tissues” (Gershoni 2017 at 2-3.) Some examples of tissues identified by these authors that have SDE genes include breast mammary tissue, skeletal muscle, skin, thyroid gland, pituitary gland, subcutaneous adipose, lung, and heart left ventricle. Based on these observations the authors state “As expected, Y-linked genes that are normally carried only by men show SDE in many tissues” (Gershoni 2017 at 3.) As stated by Heydari et al. (2022, at 1), “Y chromosome harbors male-specific genes, which either solely or in cooperation with their X-counterpart, and independent or in conjunction with sex hormones have a considerable impact on basic physiology and disease mechanisms in most or all tissues development.”

6. In a review of 56 articles on the topic of sex-based differences in skeletal muscle, Haizlip et al., (2015) state that “More than 3,000 genes have been

identified as being differentially expressed between male and female skeletal muscle.” (Haizlip 2015 at 30.) Furthermore, the authors state that “Overall, evidence to date suggests that skeletal muscle fiber-type composition is dependent on species, anatomical location/function, and sex” (Haizlip 2015 at 30.) The differences in genetic expression between males and females influence the skeletal muscle fiber composition (i.e. fast twitch and fast twitch sub-type and slow twitch), the skeletal muscle fiber size, the muscle contractile rate, and other aspects of muscle function that influence athletic performance. As the authors review the differences in skeletal muscle between males and females they conclude, “Additionally, all of the fibers measured in men have significantly larger cross-sectional areas (CSA) compared with women.” (Haizlip 2015 at 31.) The authors also explore the effects of thyroid hormone, estrogen, and testosterone on gene expression and skeletal muscle function in males and females. One major conclusion by the authors is that “The complexity of skeletal muscle and the role of sex adding to that complexity cannot be overlooked.” (Haizlip 2015 at 37.) The evaluation of SDE in protein coding genes helps illustrate that the differences between men and women are intrinsically part of the chromosomal and genetic makeup of humans which can influence many tissues that are inherent to the athletic competitive advantages of men compared to women.

II. Biological men, or adolescent boys, have large, well-documented performance advantages over women and adolescent girls in almost all athletic contests.

7. It should scarcely be necessary to invoke scientific experts to “prove” that men are on average larger, stronger, and faster than women. All of us, along with our siblings and our peers and perhaps our children, have passed through puberty, and we have watched that differentiation between the sexes occur. This is common human experience and knowledge.

8. Nevertheless, these differences have been extensively studied and measured. I cited many of these studies in the first paper on this topic that I prepared, which was submitted in litigation in January 2020. Since then, in light of current controversies, several authors have compiled valuable collections or reviews of data extensively documenting this objective fact about the human species, as manifest in almost all sports, each of which I have reviewed and found informative. These include Coleman (2020), Hilton & Lundberg (2021), World Rugby (2020), Harper (2021), Hamilton (2021), and a “Briefing Book” prepared by the Women’s Sports Policy Working Group (2021). The important paper by Handelsman et al. (2018) also gathers scientific evidence of the systematic and large male athletic advantage.

9. These papers and many others document that men, adolescent boys, and prepubertal male children, substantially outperform comparably aged women,

adolescent girls and prepubertal female children, in competitions involving running speed, swimming speed, cycling speed, jumping height, jumping distance, and strength (to name a few, but not all, of the performance differences). As I discuss later, it is now clear that these performance advantages for men, adolescent boys, and prepubertal male children, are inherent to the biological differences between the sexes.

10. In fact, I am not aware of any scientific evidence today that disproves that after puberty men possess large advantages in athletic performance over women—so large that they are generally insurmountable for comparably gifted and trained athletes at every level (i.e. (a) elite, (b) collegiate, (c) scholastic, and (d) recreational competition). And I am not aware of any scientific evidence today that disproves that these measured performance advantages are at least largely the result of physiological differences between men and women which have been measured and are reasonably well understood.

11. My use of the term “advantage” in this paper must not be read to imply any normative judgment. The adult female physique is simply different from the adult male physique. Obviously, it is optimized in important respects for the difficult task of childbearing. On average, women require far fewer calories for healthy survival. Evolutionary biologists can and do theorize about the survival value or “advantages” provided by these and other distinctive characteristics of the female physique, but I will leave that to the evolutionary biologists. I use “advantage” to refer merely to performance advantages in athletic competitions.

12. I find in the literature a widespread consensus that the large performance and physiological advantages possessed by males—rather than social considerations or considerations of identity—are precisely the *reason* that most athletic competitions are separated by sex, with women treated as a “protected class.” To cite only a few statements accepting this as the justification:

- Handelsman et al. (2018) wrote, “Virtually all elite sports are segregated into male and female competitions. The main justification is to allow women a chance to win, as women have major disadvantages against men who are, on average, taller, stronger, and faster and have greater endurance due to their larger, stronger, muscles and bones as well as a higher circulating hemoglobin level.” (803)
- Millard-Stafford et al. (2018) wrote “Current evidence suggests that women will not swim or run as fast as men in Olympic events, which speaks against eliminating sex segregation in these individual sports” (530) “Given the historical context (2% narrowing in swimming over 44 y), a reasonable assumption might be that no more than 2% of the

current performance gap could still potentially be attributed to sociocultural influences.”, (533) and “Performance gaps between US men and women stabilized within less than a decade after federal legislation provided equal opportunities for female participation, but only modestly closed the overall gap in Olympic swimming by 2% (5% in running).” (533) Dr. Millard-Stafford, a full professor at Georgia Tech, holds a Ph.D. in Exercise Physiology and is a past President of the American College of Sports Medicine.

- In 2021, Hilton et al. wrote, “most sports have a female category the purpose of which is the protection of both fairness and, in some sports, safety/welfare of athletes who do not benefit from the physiological changes induced by male levels of testosterone from puberty onwards.” (204)
- In 2020 the Swiss High Court (“Tribunal Fédéral”) observed that “in most sports . . . women and men compete in two separate categories, because the latter possess natural advantages in terms of physiology.”¹
- The members of the Women’s Sports Policy Working Group wrote that “If sports were not sex-segregated, female athletes would rarely be seen in finals or on victory podiums,” and that “We have separate sex sport and eligibility criteria based on biological sex because this is the only way we can assure that female athletes have the same opportunities as male athletes not only to participate but to win in competitive sport. . . . If we did not separate athletes on the basis of biological sex—if we used any other physical criteria—we would never see females in finals or on podiums.” (WSPWG Briefing Book 2021 at 5, 20.)
- In 2020, the World Rugby organization stated that “the women's category exists to ensure protection, safety and equality for those who do not benefit from the biological advantage created by these biological performance attributes.” (World Rugby Transgender Women Guidelines 2020.)
- In 2021 Harper et al. stated “...the small decrease in strength in transwomen after 12–36 months of GAHT [Gender Affirming Hormone Therapy] suggests that transwomen likely retain a strength advantage

¹ “dans la plupart des sports . . . les femmes et les hommes concourent dans deux catégories séparées, ces derniers étant naturellement avantagés du point de vue physique.” Tribunal Fédéral decision of August 25, 2020, Case 4A_248/2019, 4A_398/2019, at §9.8.3.3.

over cisgender women.” (7) and “...observations in trained transgender individuals are consistent with the findings of the current review in untrained transgender individuals, whereby 30 months of GAHT may be sufficient to attenuate some, but not all, influencing factors associated with muscular endurance and performance.” (8)

- Hamilton et al. (2021), in a consensus statement for the International Federation of Sports Medicine (FIMS) concluded that “Transwomen have the right to compete in sports. However, cisgender women have the right to compete in a protected category.” (1409)

13. While the sources I mention above gather more extensive scientific evidence of this uncontroversial truth, I provide here a brief summary of representative facts concerning the male advantage in athletic performance.

A. Men are stronger.

14. Males exhibit greater strength throughout the body. Both Handelsman et al. (2018) and Hilton & Lundberg (2021) have gathered multiple literature references that document this fact in various muscle groups.

15. Men have in the neighborhood of 60%-100% greater **arm strength** than women. (Handelsman 2018 at 812.)² One study of elbow flexion strength (basically, bringing the fist up towards the shoulder) in a large sample of men and women found that men exhibited 109% greater isometric strength, and 89% higher strength in a single repetition. (Hilton 2021 at 204, summarizing Hubal (2005) at Table 2.)

16. **Grip strength** is often used as a useful proxy for strength more generally. In one study, men showed on average 57% greater grip strength than women. (Bohannon 2019.) A wider meta-analysis of multiple grip-strength studies not limited to athletic populations found that 18- and 19-year-old males exhibited in

² Handelsman expresses this as women having 50% to 60% of the “upper limb” strength of men. Handelsman cites Sale, *Neuromuscular function*, for this figure and the “lower limb” strength figure. Knox et al., *Transwomen in elite sport* (2018) are probably confusing the correct way to state percentages when they state that “differences lead to decreased trunk and lower body strength by 64% and 72% respectively, in women” (397): interpreted literally, this would imply that men have **almost 4x as much** lower body strength as do women.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

the neighborhood of 2/3 greater grip strength than females. (Handelsman 2017 Figure 3, summarizing Silverman 2011 Table 1.)³

17. In an evaluation of maximal isometric handgrip strength in 1,654 healthy men, 533 healthy women aged 20-25 years and 60 “highly trained elite female athletes from sports known to require high hand-grip forces (judo, handball),” Leyk et al. (2007) observed that, “The results of female national elite athletes even indicate that the strength level attainable by extremely high training will rarely surpass the 50th percentile of untrained or not specifically trained men.” (Leyk 2007 at 415.)

18. Men have in the neighborhood of 25%-60% greater **leg strength** than women. (Handelsman 2018 at 812.) In another measure, men exhibit 54% greater knee extension torque and this male leg strength advantage is consistent across the lifespan. (Neder 1999 at 120-121.)

19. When male and female Olympic weightlifters of the same body weight are compared, the top males lift weights between 30% and 40% greater than the females of the same body weight. But when top male and female performances are compared in powerlifting, without imposing any artificial limitations on bodyweight, the male record is 65% higher than the female record. (Hilton 2021 at 203.)

20. In another measure that combines many muscle groups as well as weight and speed, moderately trained males generated 162% greater punching power than females even though men do not possess this large an advantage in any single bio-mechanical variable. (Morris 2020.) This objective reality was subjectively summed up by women’s mixed-martial arts fighter Tamikka Brents, who suffered significant facial injuries when she fought against a biological male who identified as female and fought under the name of Fallon Fox. Describing the experience, Brents said:

“I’ve fought a lot of women and have never felt the strength that I felt in a fight as I did that night. I can’t answer whether it’s because she was born a man or not because I’m not a doctor. I can only say, I’ve never felt so overpowered ever in my life, and I am an abnormally strong female in my own right.”⁴

³ Citing Silverman, *The secular trend for grip strength in Canada and the United States*, J. Ports Sci. 29:599-606 (2011).

⁴ <http://whoatv.com/exclusive-fallon-foxs-latest-opponent-opens-up-to-whoatv/> (last accessed October 5, 2021).

B. Men run faster.

21. Many scholars have detailed the wide performance advantages enjoyed by men in running speed. One can come at this reality from a variety of angles.


22. Multiple authors report a male speed advantage in the neighborhood of 10%-13% in a variety of events, with a variety of study populations. Handelsman et al. 2018 at 813 and Handelsman 2017 at 70 both report a male advantage of about 10% by age 17. Thibault et al. 2010 at 217 similarly reported a stable 10% performance advantage across multiple events at the Olympic level. Tønnessen et al. (2015 at 1-2) surveyed the data and found a consistent male advantage of 10%-12% in running events after the completion of puberty. They document this for both short sprints and longer distances. One group of authors found that the male advantage increased dramatically in ultra-long-distance competition (Lepers & Knechtle 2013.)

23. A great deal of current interest has been focused on track events. It is worth noting that a recent analysis of publicly available sports federation and tournament records found that men enjoy the *least* advantage in running events, as compared to a range of other events and metrics, including jumping, pole vaulting, tennis serve speed, golf drives, baseball pitching speed, and weightlifting. (Hilton 2021 at 201-202.) Nevertheless, as any serious runner will recognize, the approximately 10% male advantage in running is an overwhelming difference. Dr. Hilton calculates that “approximately 10,000 males have personal best times that are faster than the current Olympic 100m female champion.” (Hilton 2021 at 204.) Professors Doriane Coleman, Jeff Wald, Wickliffe Shreve, and Richard Clark dramatically illustrated this by compiling the data and creating the figure below (last accessed on February 10, 2022, at <https://bit.ly/35yOyS4>), which shows that the *lifetime best performances* of three female Olympic champions in the 400m event—including Team USA’s Sanya Richards-Ross and Allyson Felix—would not match the performances of “literally thousands of boys and men, including thousands who would be considered second tier in the men’s category” *just in 2017 alone*: (data were drawn from the International Association of Athletics Federations (IAAF) website which provides complete, worldwide results for individuals and events, including on an annual and an all-time basis).

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Comparing the Best Elite Females to Boys and Men:
Personal Bests for 3 Female Gold Medalists versus 2017 Performances by Boys and Men



G. Brown

Expert Report, B.P.J. v. WV BOE et al.

24. Professor Coleman and her colleague Wicklyffe Shreve also created the table below (last accessed on February 10, 2022, at <https://bit.ly/37E1s2X>), which “compares the number of men—males over 18—competing in events reported to the International Association of Athletics Federation whose results in each event in 2017 would have ranked them above the very best elite woman that year.”

TABLE 2 – World’s Best Woman v. Number of Men Outperforming			
Event	Best Women’s Result	Best Men’s Result	# of Men Outperforming
100 Meters	10.71	9.69	2,474
200 Meters	21.77	19.77	2,920
400 Meters	49.46	43.62	4,341
800 Meters	1:55.16*	1:43.10	3,992+
1500 Meters	3:56.14	3:28.80	3,216+
3000 Meters	8:23.14	7:28.73	1307+
5000 Meters	14:18.37	12:55.23	1,243
High Jump	2.06 meters	2.40 meters	777
Pole Vault	4.91 meters	6.00 meters	684
Long Jump	7.13 meters	8.65 meters	1,652
Triple Jump	14.96 meters	18.11 meters	969

25. The male advantage becomes insuperable well before the developmental changes of puberty are complete. Dr. Hilton documents that even “schoolboys”—defined as age 15 and under—have beaten the female world records in running, jumping, and throwing events. (Hilton 2021 at 204.)

26. Similarly, Coleman and Shreve created the table below (last accessed on February 10, 2022, at <https://bit.ly/37E1s2X>), which “compares the number of boys—males under the age of 18—whose results in each event in 2017 would rank them above the single very best elite [adult] woman that year.” data were drawn from the International Association of Athletics Federations (IAAF) website

TABLE 1 – World’s Best Woman v. Under 18 Boys			
Event	Best Women’s Result	Best Boys’ Result	# of Boys Outperforming
100 Meters	10.71	10.15	124 ⁺
200 Meters	21.77	20.51	182
400 Meters	49.46	45.38	285
800 Meters	1:55.16*	1:46.3	201+
1500 Meters	3:56.14	3:37.43	101+
3000 Meters	8:23.14	7:38.90	30
5000 Meters	14:18.37	12:55.58	15
High Jump	2.06 meters	2.25 meters	28
Pole Vault	4.91 meters	5.31 meters	10
Long Jump	7.13 meters	7.88 meters	74
Triple Jump	14.96 meters	17.30 meters	47

27. In an analysis I have performed of running events (consisting of the 100 m, 200 m, 400 m, 800 m, 1500 m, 5000 m, and 10000 m) in the Division 1, Division 2, and Division 3 NCAA Outdoor track championships for the years of 2010-2019, the average performance across all events of the 1st place man was 14.1% faster than the 1st place woman, with the smallest difference being a 10.2% advantage for men in the Division 1 100 m race. The average 8th place man across all events (the last place to earn the title of All American) was 11.2% faster than 1st place woman, with the smallest difference being a 6.5% advantage for men in the Division 1 100 m race. (Brown et al. Unpublished observations, to be presented at the 2022 Annual Meeting of the American College of Sports Medicine.)

28. Athletic.net® is an internet-based resource providing “results, team, and event management tools to help coaches and athletes thrive.” Among the resources available on Athletic.net are event records that can be searched by nationally or by state age group, school grade, and state. Higerd (2021) in an evaluation of high school track running performance records from five states (CA, FL, MN, NY, WA), over three years (2017 – 2019) observed that males were 14.38% faster than females in the 100M (at 99), 16.17% faster in the 200M (at 100), 17.62% faster in the 400M (at 102), 17.96% faster in the 800M (at 103), 17.81% faster in the 1600M (at 105), and 16.83% faster in the 3200M (at 106).

C. Men jump higher and farther.

29. Jumping involves both leg strength and speed as positive factors, with body weight of course a factor working against jump height. Despite their substantially greater body weight, males enjoy an even greater advantage in jumping than in running. Handelsman 2018 at 813, looking at youth and young adults, and Thibault 2010 at 217, looking at Olympic performances, both found male advantages in the range of 15%-20%. See also Tønnessen 2015 (approximately 19%); Handelsman 2017 (19%); Hilton 2021 at 201 (18%). Looking at the vertical jump called for in volleyball, research on elite volleyball players found that males jumped on average 50% higher during an “attack” at the net than did females. (Sattler 2015; see also Hilton 2021 at 203 (33% higher vertical jump).)

30. Higerd (2021) in an evaluation of high school high jump performance available through the track and field database athletic.net®, which included five states (CA, FL, MN, NY, WA), over three years (2017 – 2019) (at 82) observed that in 23,390 females and 26,843 males, females jumped an average of 1.35 m and males jumped an average of 1.62 m, for an 18.18% performance advantage for males (at 96). In an evaluation of long jump performance in 45,705 high school females and 54,506 high school males the females jumped an average of 4.08 m and males jumped an average of 5.20 m, for a 24.14% performance advantage for males (at 97).

31. The combined male advantage of body height and jump height means, for example, that a total of seven women in the WNBA have ever dunked a basketball in the regulation 10 foot hoop,⁵ while the ability to dunk appears to be almost universal among NBA players: “Since the 1996–97 season (the earliest data is available from Basketball-Reference.com), 1,801 different [NBA] players have combined for 210,842 regular-season dunks, and 1,259 out of 1,367 players (or 92%) who have played at least 1,000 minutes have dunked at least once.”⁶

D. Men throw, hit, and kick faster and farther.

32. Strength, arm-length, and speed combine to give men a large advantage over women in throwing. This has been measured in a number of studies.

33. One study of elite male and female baseball pitchers showed that men throw baseballs 35% faster than women—81 miles/hour for men vs. 60 miles/hour for women. (Chu 2009.) By age 12, “boys’ throwing velocity is already between 3.5 and 4 standard deviation units higher than the girls’.” (Thomas 1985 at 276.) By age seventeen, the *average* male can throw a ball farther than 99% of seventeen-year-old females. (Lombardo 2018; Chu 2009; Thomas 1985 at 268.) Looking at publicly available data, Hilton & Lundberg found that in both baseball pitching and the field hockey “drag flick,” the *record* ball speeds achieved by males are more than 50% higher than those achieved by females. (Hilton 2021 at 202-203.)

34. Men achieve serve speeds in tennis more that 15% faster than women; and likewise in golf achieve ball speeds off the tee more than 15% faster than women. (Hilton 2021 at 202.)

35. Males are able to throw a javelin more than 30% farther than females. (Lombardo 2018 Table 2; Hilton 2021 at 203.)

36. Men serve and spike volleyballs with higher velocity than women, with a performance advantage in the range of 29-34%. (Hilton 2021 at 204 Fig. 1.)

37. Men are also able to kick balls harder and faster. A study comparing collegiate soccer players found that males kick the ball with an average 20% greater velocity than females. (Sakamoto 2014.)

⁵ https://www.espn.com/wnba/story/_/id/32258450/2021-wnba-playoffs-brittney-griner-owns-wnba-dunking-record-coming-more.

⁶ <https://www.si.com/nba/2021/02/22/nba-non-dunkers-patty-mills-tj-mcconnell-steve-novak-daily-cover>

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

E. Males exhibit faster reaction times.

38. Interestingly, men enjoy an additional advantage over women in reaction time—an attribute not obviously related to strength or metabolism (e.g. VO_2max). “Reaction time in sports is crucial in both simple situations such as the gun shot in sprinting and complex situations when a choice is required. In many team sports this is the foundation for tactical advantages which may eventually determine the outcome of a game.” (Dogan 2009 at 92.) “Reaction times can be an important determinant of success in the 100m sprint, where medals are often decided by hundredths or even thousandths of a second.” (Tønnessen 2013 at 885.)

39. The existence of a sex-linked difference in reaction times is consistent over a wide range of ages and athletic abilities. (Dykiert 2012.) Even by the age of 4 or 5, in a ruler-drop test, males have been shown to exhibit 4% to 6% faster reaction times than females. (Latorre-Roman 2018.) In high school athletes taking a common baseline “ImPACT” test, males showed 3% faster reaction times than females. (Mormile 2018.) Researchers have found a 6% male advantage in reaction times of both first-year medical students (Jain 2015) and world-class sprinters (Tønnessen 2013).

40. Most studies of reaction times use computerized tests which ask participants to hit a button on a keyboard or to say something in response to a stimulus. One study on NCAA athletes measured “reaction time” by a criterion perhaps more closely related to athletic performance—that is, how fast athletes covered 3.3 meters after a starting signal. Males covered the 3.3 meters 10% faster than females in response to a visual stimulus, and 16% faster than females in response to an auditory stimulus. (Spierer 2010.)

41. Researchers have speculated that sex-linked differences in brain structure, as well as estrogen receptors in the brain, may be the source of the observed male advantage in reaction times, but at present this remains a matter of speculation and hypothesis. (Mormile at 19; Spierer at 962.)

III. Men have large measured physiological differences compared to women which demonstrably or likely explain their performance advantages.

42. No single physiological characteristic alone accounts for all or any one of the measured advantages that men enjoy in athletic performance. However, scientists have identified and measured a number of physiological factors that contribute to superior male performance.

A. Men are taller and heavier than women

43. In some sports, such as basketball and volleyball, height itself provides competitive advantage. While some women are taller than some men, based on data from 20 countries in North America, Europe, East Asia, and Australia, the 50th percentile for body height for women is 164.7 cm (5 ft 5 inches) and the 50th percentile for body height for men is 178.4 cm (5 ft 10 inches). Helping to illustrate the inherent height difference between men and women, from the same data analysis, the 95th percentile for body height for women is 178.9 cm (5 feet 10.43 inches), which is only 0.5 cm taller than the 50th percentile for men (178.4 cm; 5 feet 10.24 inches), while the 95th percentile for body height for men is 193.6 cm (6 feet 4.22 inches). (Roser 2013.)

44. To look at a specific athletic population, an evaluation of NCAA Division 1 basketball players compared 68 male guards and 59 male forwards to 105 female guards and 91 female forwards, and found that on average the male guards were 187.4 ± 7.0 cm tall and weighed 85.2 ± 7.4 kg while the female guards were 171.6 ± 5.0 cm tall and weighed 68.0 ± 7.4 kg. The male forwards were 201.7 ± 4.0 cm tall and weighed 105.3 ± 5.9 kg while the female forwards were 183.5 ± 4.4 cm tall and weighed 82.2 ± 12.5 kg. (Fields 2018 at 3.)

B. Males have larger and longer bones, stronger bones, and different bone configuration.

45. Obviously, males on average have longer bones. “Sex differences in height have been the most thoroughly investigated measure of bone size, as adult height is a stable, easily quantified measure in large population samples. Extensive twin studies show that adult height is highly heritable with predominantly additive genetic effects that diverge in a sex-specific manner from the age of puberty onwards.” (Handelsman 2018 at 818.) “Pubertal testosterone exposure leads to an ultimate average greater height in men of 12–15 centimeters, larger bones, greater muscle mass, increased strength and higher hemoglobin levels.” (Gooren 2011 at 653.)

46. “Men have distinctively greater bone size, strength, and density than do women of the same age. As with muscle, sex differences in bone are absent prior to puberty but then accrue progressively from the onset of male puberty due to the sex difference in exposure to adult male circulating testosterone concentrations.” (Handelsman 2018 at 818.)

47. “[O]n average men are 7% to 8% taller with longer, denser, and stronger bones, whereas women have shorter humerus and femur cross-sectional

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

areas being 65% to 75% and 85%, respectively, those of men.” (Handelsman 2018 at 818.)

48. Greater height, leg, and arm length themselves provide obvious advantages in several sports. But male bone geometry also provides less obvious advantages. “The major effects of men’s larger and stronger bones would be manifest via their taller stature as well as the larger fulcrum with greater leverage for muscular limb power exerted in jumping, throwing, or other explosive power activities.” (Handelsman 2018 at 818.)

49. Male advantage in bone size is not limited to length, as larger bones provide the mechanical framework for larger muscle mass. “From puberty onwards, men have, on average, 10% more bone providing more surface area. The larger surface area of bone accommodates more skeletal muscle so, for example, men have broader shoulders allowing more muscle to build. This translates into 44% less upper body strength for women, providing men an advantage for sports like boxing, weightlifting and skiing. In similar fashion, muscle mass differences lead to decreased trunk and lower body strength by 64% and 72%, respectively in women. These differences in body strength can have a significant impact on athletic performance, and largely underwrite the significant differences in world record times and distances set by men and women.” (Knox 2019 at 397.)

50. Meanwhile, distinctive aspects of the female pelvis geometry cut against athletic performance. “[T]he widening of the female pelvis during puberty, balancing the evolutionary demands of obstetrics and locomotion, retards the improvement in female physical performance.” (Handelsman 2018 at 818.) “[T]he major female hormones, oestrogens, can have effects that disadvantage female athletic performance. For example, women have a wider pelvis changing the hip structure significantly between the sexes. Pelvis shape is established during puberty and is driven by oestrogen. The different angles resulting from the female pelvis leads to decreased joint rotation and muscle recruitment ultimately making them slower.” (Knox 2019 at 397.)

51. There are even sex-based differences in foot size and shape. Wunderlich & Cavanaugh (2001) observed that a “foot length of 257 mm represents a value that is ... approximately the 20th percentile men’s foot lengths and the 80th percentile women’s foot lengths.” (607) and “For a man and a woman, both with statures of 170 cm (5 feet 7 inches), the man would have a foot that was approximately 5 mm longer and 2 mm wider than the woman.” (608). Based on these, and other analyses, they conclude that “female feet and legs are not simply scaled-down versions of male feet but rather differ in a number of shape characteristics, particularly at the arch, the lateral side of the foot, the first toe, and the ball of the foot.” (605) Further, Fessler et al. (2005) observed that “female foot length is consistently smaller than male foot length” (44) and concludes that

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

“proportionate foot length is smaller in women” (51) with an overall conclusion that “Our analyses of genetically disparate populations reveal a clear pattern of sexual dimorphism, with women consistently having smaller feet proportionate to stature than men.” (53)

52. Beyond simple performance, the greater density and strength of male bones provide higher protection against stresses associated with extreme physical effort: “[S]tress fractures in athletes, mostly involving the legs, are more frequent in females, with the male protection attributable to their larger and thicker bones.” (Handelsman 2018 at 818.)

C. Males have much larger muscle mass.

53. The fact that, on average, men have substantially larger muscles than women is as well known to common observation as men’s greater height. But the male advantage in muscle size has also been extensively measured. The differential is large.

54. “On average, women have 50% to 60% of men’s upper arm muscle cross-sectional area and 65% to 70% of men’s thigh muscle cross-sectional area, and women have 50% to 60% of men’s upper limb strength and 60% to 80% of men’s leg strength. Young men have on average a skeletal muscle mass of >12 kg greater than age-matched women at any given body weight.” (Handelsman 2018 at 812. See also Gooren 2011 at 653, Thibault 2010 at 214.)

55. “There is convincing evidence that the sex differences in muscle mass and strength are sufficient to account for the increased strength and aerobic performance of men compared with women and is in keeping with the differences in world records between the sexes.” (Handelsman 2018 at 816.)

56. Once again, looking at specific and comparable populations of athletes, an evaluation of NCAA Division 1 basketball players consisting of 68 male guards and 59 male forwards, compared to 105 female guards and 91 female forwards, reported that on average the male guards had 77.7 ± 6.4 kg of fat free mass and 7.4 ± 3.1 kg fat mass while the female guards had 54.6 ± 4.4 kg fat free mass and 13.4 ± 5.4 kg fat mass. The male forwards had 89.5 ± 5.9 kg fat free mass and 15.9 ± 5.6 kg fat mass while the female forwards had 61.8 ± 5.9 kg fat free mass and 20.5 ± 7.7 kg fat mass. (Fields 2018 at 3.)

D. Females have a larger proportion of body fat.

57. While women have smaller muscles, they have proportionately more body fat, in general a negative for athletic performance. “Oestrogens also affect body

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

composition by influencing fat deposition. Women, on average, have higher percentage body fat, and this holds true even for highly trained healthy athletes (men 5%–10%, women 8%–15%). Fat is needed in women for normal reproduction and fertility, but it is not performance-enhancing. This means men with higher muscle mass and less body fat will normally be stronger kilogram for kilogram than women.” (Knox 2019 at 397.)

58. “[E]lite females have more (<13 vs. <5 %) body fat than males. Indeed, much of the difference in [maximal oxygen uptake] between males and females disappears when it is expressed relative to lean body mass. . . . Males possess on average 7–9 % less percent body fat than females.” (Lepers 2013 at 853.)

59. Knox et al. observe that both female pelvis shape and female body fat levels “disadvantage female athletes in sports in which speed, strength and recovery are important,” (Knox 2019 at 397), while Tønnessen et al. describe the “ratio between muscular power and total body mass” as “critical” for athletic performance. (Tønnessen 2015 at 7.)

E. Males are able to metabolize and release energy to muscles at a higher rate due to larger heart and lung size, and higher hemoglobin concentrations.

60. While advantages in bone size, muscle size, and body fat are easily perceived and understood by laymen, scientists also measure and explain the male athletic advantage at a more abstract level through measurements of metabolism, or the ability to deliver energy to muscles throughout the body.

61. Energy release at the muscles depends centrally on the body’s ability to deliver oxygen to the muscles, where it is essential to the complex chain of biochemical reactions that make energy available to power muscle fibers. Men have multiple distinctive physiological attributes that together give them a large advantage in oxygen delivery.

62. Oxygen is taken into the blood in the lungs. Men have greater capability to take in oxygen for multiple reasons. “[L]ung capacity [is] larger in men because of a lower diaphragm placement due to Y-chromosome genetic determinants.” (Knox 2019 at 397.) Supporting larger lung capacity, men have “greater cross-sectional area of the trachea”; that is, they can simply move more air in and out of their lungs in a given time. (Hilton 2021 at 201.)

63. More, male lungs provide superior oxygen exchange even for a given volume: “The greater lung volume is complemented by testosterone-driven **enhanced alveolar multiplication** rate during the early years of life. Oxygen exchange takes place between the air we breathe and the bloodstream at the alveoli,

so more alveoli allows more oxygen to pass into the bloodstream. Therefore, the greater lung capacity allows more air to be inhaled with each breath. This is coupled with an improved uptake system allowing men to absorb more oxygen.” (Knox 2019 at 397.)

64. “Once in the blood, oxygen is carried by haemoglobin. **Haemoglobin concentrations** are directly modulated by testosterone so men have higher levels and can carry more oxygen than women.” (Knox 2019 at 397.) “It is well known that levels of circulating hemoglobin are androgen-dependent and consequently higher in men than in women by 12% on average.... Increasing the amount of hemoglobin in the blood has the biological effect of increasing oxygen transport from lungs to tissues, where the increased availability of oxygen enhances aerobic energy expenditure.” (Handelsman 2018 at 816.) (See also Lepers 2013 at 853; Handelsman 2017 at 71.) “It may be estimated that as a result the average maximal oxygen transfer will be ~10% greater in men than in women, which has a direct impact on their respective athletic capacities.” (Handelsman 2018 at 816.)

65. But the male metabolic advantage is further multiplied by the fact that men are also able to **circulate more blood per second** than are women. “Oxygenated blood is pumped to the active skeletal muscle by the heart. The left ventricle chamber of the heart is the reservoir from which blood is pumped to the body. The larger the left ventricle, the more blood it can hold, and therefore, the more blood can be pumped to the body with each heartbeat, a physiological parameter called ‘stroke volume’. The female heart size is, on average, 85% that of a male resulting in the stroke volume of women being around 33% less.” (Knox 2018 at 397.) Hilton cites different studies that make the same finding, reporting that men on average can pump 30% more blood through their circulatory system per minute (“cardiac output”) than can women. (Hilton 2021 at 202.)

66. Finally, at the cell where the energy release is needed, men appear to have yet another advantage. “Additionally, there is experimental evidence that testosterone increases . . . **mitochondrial biogenesis**, myoglobin expression, and IGF-1 content, which may augment energetic and power generation of skeletal muscular activity.” (Handelsman 2018 at 811.)

67. “Putting all of this together, men have a much more efficient cardiovascular and respiratory system.” (Knox 2019 at 397.) A widely accepted measurement that reflects the combined effects of all these respiratory, cardiovascular, and metabolic advantages is referred to as “ $\dot{V}O_{2\max}$,” which refers to the maximum rate at which an individual can consume oxygen during aerobic

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

exercise.⁷ Looking at 11 separate studies, including both trained and untrained individuals, Pate et al. concluded that men have a 50% higher VO_2max than women on average, and a 25% higher VO_2max in relation to body weight. (Pate 1984 at 92. See also Hilton 2021 at 202.)

IV. The role of testosterone in the development of male advantages in athletic performance.

68. The following tables of reference ranges for circulating testosterone in males and females are presented to help provide context for some of the subsequent information regarding athletic performance and physical fitness in children, youth, and adults, and regarding testosterone suppression in transwomen and athletic regulations. These data were obtained from the Mayo Clinic Laboratories (available at <https://www.mayocliniclabs.com/test-catalog/overview/83686#Clinical-and-Interpretive>, accessed January 14, 2022).

Reference ranges for serum testosterone concentrations in males and females.

Age	Males	Females
0 – 5 months	2.6 – 13.9 nmol/l	0.7 – 2.8 nmol/l
6 months – 9 years	0.2 – 0.7 nmol/l	0.2 – 0.7 nmol/l
10 – 11 years	0.2 – 4.5 nmol/l	0.2 – 1.5 nmol/l
12 -13 years	0.2 – 27.7 nmol/l	0.2 – 2.6 nmol/l
14 years	0.2 – 41.6 nmol/l	0.2 – 2.6 nmol/l
15 – 16 years	3.5 – 41.6 nmol/l	0.2 – 2.6 nmol/l
17 – 18 years	10.4 – 41.6 nmol/l	0.7 – 2.6 nmol/l
19 years and older	8.3 – 32.9 nmol/l	0.3 – 2.1 nmol/l

Please note that testosterone concentrations are sometimes expressed in units of ng/dl, and 1 nmol/l = 28.85 ng/dl.

69. Tanner Stages can be used to help evaluate the onset and progression of puberty and may be more helpful in evaluating normal testosterone concentrations than age in adolescents. “Puberty onset (transition from Tanner stage I to Tanner stage II) occurs for boys at a median age of 11.5 years and for girls

⁷ VO_2max is “based on hemoglobin concentration, total blood volume, maximal stroke volume, cardiac size/mass/compliance, skeletal muscle blood flow, capillary density, and mitochondrial content.” International Statement, *The Role of Testosterone in Athletic Performance* (January 2019), available at https://law.duke.edu/sites/default/files/centers/sportslaw/Experts_T_Statement_2019.pdf.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

at a median age of 10.5 years. . . . Progression through Tanner stages is variable. Tanner stage V (young adult) should be reached by age 18.”

(<https://www.mayocliniclabs.com/test-catalog/overview/83686#Clinical-and-Interpretive>, accessed January 14, 2022).

Reference Ranges for serum testosterone concentrations by Tanner stage

Tanner Stage	Males	Females
I (prepubertal)	0.2 – 0.7 nmol/l	0.7 – 0.7 nmol/l
II	0.3 – 2.3 nmol/l	0.2 – 1.6 nmol/l
III	0.9 – 27.7 nmol/l	0.6 – 2.6 nmol/l
IV	2.9 – 41.6 nmol/l	0.7 – 2.6 nmol/l
V (young adult)	10.4 – 32.9 nmol/l	0.4 – 2.1 nmol/l

70. Senefeld et al. (2020 at 99) state that “Data on testosterone levels in children and adolescents segregated by sex are scarce and based on convenience samples or assays with limited sensitivity and accuracy.” They therefore “analyzed the timing of the onset and magnitude of the divergence in testosterone in youths aged 6 to 20 years by sex using a highly accurate assay” (isotope dilution liquid chromatography tandem mass spectrometry). Senefeld observed a significant difference beginning at age 11, which is to say about fifth grade.

Serum testosterone concentrations (nmol/L) in youths aged 6 to 20 years measured using isotope dilution liquid chromatography tandem mass spectrometry (Senefeld et al. ,2020, at 99)

Age (y)	Boys			Girls		
	5th	50th	95th	5th	50th	95th
6	0.0	0.1	0.2	0.0	0.1	0.2
7	0.0	0.1	0.2	0.0	0.1	0.3
8	0.0	0.1	0.3	0.0	0.1	0.3
9	0.0	0.1	0.3	0.1	0.2	0.6
10	0.1	0.2	2.6	0.1	0.3	0.9
11	0.1	0.5	11.3	0.2	0.5	1.3
12	0.3	3.6	17.2	0.2	0.7	1.4
13	0.6	9.2	21.5	0.3	0.8	1.5
14	2.2	11.9	24.2	0.3	0.8	1.6
15	4.9	13.2	25.8	0.4	0.8	1.8
16	5.2	14.9	24.1	0.4	0.9	2.0
17	7.6	15.4	27.0	0.5	1.0	2.0
18	9.2	16.3	25.5	0.4	0.9	2.1
19	8.1	17.2	27.9	0.4	0.9	2.3
20	6.5	17.9	29.9	0.4	1.0	3.4

A. Boys exhibit advantages in athletic performance even before puberty.

71. It is often said or assumed that boys enjoy no significant athletic advantage over girls before puberty. However, this is not true. Writing in their seminal work on the physiology of elite young female athletes, McManus and Armstrong (2011) reviewed the differences between boys and girls regarding bone density, body composition, cardiovascular function, metabolic function, and other physiologic factors that can influence athletic performance. They stated, “At birth, boys tend to have a greater lean mass than girls. This difference remains small but detectable throughout childhood with about a 10% greater lean mass in boys than girls prior to puberty.” (28) “Sexual dimorphism underlies much of the physiologic response to exercise,” and most importantly these authors concluded that, “Young girl athletes are not simply smaller, less muscular boys.” (23)

72. Certainly, boys’ physiological and performance advantages increase rapidly from the beginning of puberty until around age 17-19. But much data and multiple studies show that significant physiological differences, and significant male athletic performance advantages in certain areas, exist before significant developmental changes associated with male puberty have occurred.

73. Starting at birth, girls have more body fat and less fat-free mass than boys. Davis et al. (2019) in an evaluation of 602 infants reported that at birth and age 5 months, infant boys have larger total body mass, body length, and fat-free mass while having lower percent body fat than infant girls. In an evaluation of 20 boys and 20 girls ages 3-8 years old, matched for age, height, and body weight Taylor et al. (Taylor 1997) reported that the “boys had significantly less fat, a lower % body fat and a higher bone-free lean tissue mass than the girls” when “expressed as a percentage of the average fat mass of the boys”, the girls’ fat mass was 52% higher than the boys “...while the bone-free lean tissue mass was 9% lower” (at 1083.) In an evaluation of 376 prepubertal [Tanner Stage 1] boys and girls, Taylor et al. (2010) observed that the boys had 21.6% more lean mass, and 13% less body fat (when expressed as percent of total body mass) than did the girls. In a review of 22 peer reviewed publications on the topic, Staiano and Katzmarzyk (2012) conclude that “... girls have more T[otal]B[ody]F[at] than boys throughout childhood and adolescence. (at 4.)

74. In the seminal textbook, *Growth, Maturation, and Physical Activity*, Malina et al. (2004) present a summary of data from Gauthier et al. (1983) which present data from “a national sample of Canadian children and youth” demonstrating that from ages 7 to 17, boys have a higher aerobic power output than do girls of the same ages when exercise intensity is measured using heart rate

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

(Malina at 242.) That is to say, that at a heart rate of 130 beats per minute, or 150, or 170, a 7 to 17 year old boy should be able to run, bike, or swim faster than a similarly aged girl.

75. Considerable data from school-based fitness testing exists showing that prepubertal boys outperform comparably aged girls in tests of muscular strength, muscular endurance, and running speed. These sex-based differences in physical fitness are relevant to the current issue of sex-based sports categories because, as stated by Lesinski et al. (2020), in an evaluation “of 703 male and female elite young athletes aged 8–18” (1) “fitness development precedes sports specialization” (2) and further observed that “males outperformed females in C[ounter]M[ovement]J[ump], D[rop]J[ump], C[hange]o[f]D[irection speed] performances and hand grip strength.” (5).

76. Tambalis et al. (2016) states that “based on a large data set comprising 424,328 test performances” (736) using standing long jump to measure lower body explosive power, sit and reach to measure flexibility, timed 30 second sit ups to measure abdominal and hip flexor muscle endurance, 10 x 5 meter shuttle run to evaluate speed and agility, and multi-stage 20 meter shuttle run test to estimate aerobic performance (738). “For each of the fitness tests, performance was better in boys compared with girls ($p < 0.001$), except for the S[it and] R[each] test ($p < 0.001$).” (739) In order to illustrate that the findings of Tambalis (2016) are not unique to children in Greece, the authors state “Our findings are in accordance with recent studies from Latvia [] Portugal [] and Australia [Catley & Tomkinson (2013)].”(744).

77. The 20-m multistage fitness test is a commonly used maximal running aerobic fitness test used in the Eurofit Physical Fitness Test Battery and the FitnessGram Physical Fitness test. It is also known as the 20-meter shuttle run test, PACER test, or beep test (among other names; this is not the same test as the shuttle run in the Presidential Fitness Test). This test involves continuous running between two lines 20 meters apart in time to recorded beeps. The participants stand behind one of the lines facing the second line and begin running when instructed by the recording. The speed at the start is quite slow. The subject continues running between the two lines, turning when signaled by the recorded beeps. After about one minute, a sound indicates an increase in speed, and the beeps will be closer together. This continues each minute (level). If the line is reached before the beep sounds, the subject must wait until the beep sounds before continuing. If the line is not reached before the beep sounds, the subject is given a warning and must continue to run to the line, then turn and try to catch up with the pace within two more 'beeps'. The subject is given a warning the first time they fail to reach the line (within 2 meters) and eliminated after the second warning.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

78. To illustrate the sex-based performance differences observed by Tambalis, I have prepared the following table showing the number of laps completed in the 20 m shuttle run for children ages 6-18 years for the low, middle, and top decile (Tambalis 2016 at 740 & 742), and have calculated the percent difference between the boys and girls using the same equation as Millard-Stafford (2018).

Performance difference between boys and girls ÷ Girls performance

Number of laps completed in the 20m shuttle run for children ages 6-18 years

Age	Male			Female			Male-Female % Difference		
	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile
6	4	14	31	4.0	12.0	26.0	0.0%	16.7%	19.2%
7	8	18	38	8.0	15.0	29.0	0.0%	20.0%	31.0%
8	9	23	47	9.0	18.0	34.0	0.0%	27.8%	38.2%
9	11	28	53	10.0	20.0	40.0	10.0%	40.0%	32.5%
10	12	31	58	11.0	23.0	43.0	9.1%	34.8%	34.9%
11	15	36	64	12.0	26.0	48.0	25.0%	38.5%	33.3%
12	15	39	69	12.0	26.0	49.0	25.0%	50.0%	40.8%
13	16	44	76	12.0	26.0	50.0	33.3%	69.2%	52.0%
14	19	50	85	12.0	26.0	50.0	58.3%	92.3%	70.0%
15	20	53	90	12.0	25.0	47.0	66.7%	112.0%	91.5%
16	20	54	90	11.0	24.0	45.0	81.8%	125.0%	100.0%
17	18	50	86	10.0	23.0	50.0	80.0%	117.4%	72.0%
18	13	48	87	8.0	23.0	39.5	62.5%	108.7%	120.3%

79. The Presidential Fitness Test was widely used in schools in the United States from the late 1950s until 2013 (when it was phased out in favor of the Presidential Youth Fitness Program and FitnessGram, both of which focus on health-related physical fitness and do not present data in percentiles). Students participating in the Presidential Fitness Test could receive “The National Physical Fitness Award” for performance equal to the 50th percentile in five areas of the fitness test, “while performance equal to the 85th percentile could receive the Presidential Physical Fitness Award.” Tables presenting the 50th and 85th percentiles for the Presidential Fitness Test for males and females ages 6 – 17, and differences in performance between males and females, for curl-ups, shuttle run, 1 mile run, push-ups, and pull-ups appear in the Appendix.

80. For both the 50th percentile (The National Physical Fitness Award) and the 85th percentile (Presidential Physical Fitness Award), with the exception of curl-ups in 6-year-old children, boys outperform girls. The difference in pull-ups for the 85th percentile for ages 7 through 17 are particularly informative with boys

outperforming girls by 100% – 1200%, highlighting the advantages in upper body strength in males.

81. A very recent literature review commissioned by the five United Kingdom governmental Sport Councils concluded that while “[i]t is often assumed that children have similar physical capacity regardless of their sex, . . . large-scale data reports on children from the age of six show that young males have significant advantage in cardiovascular endurance, muscular strength, muscular endurance, speed/agility and power tests,” although they “score lower on flexibility tests.” (UK Sports Councils’ Literature Review 2021 at 3.)

82. Hilton et al., also writing in 2021, reached the same conclusion: “An extensive review of fitness data from over 85,000 Australian children aged 9–17 years old showed that, compared with 9-year-old females, 9-year-old males were faster over short sprints (9.8%) and 1 mile (16.6%), could jump 9.5% further from a standing start (a test of explosive power), could complete 33% more push-ups in 30 [seconds] and had 13.8% stronger grip.” (Hilton 2021 at 201, summarizing the findings of Catley & Tomkinson 2013.)

83. The following data are taken from Catley & Tomkinson (2013 at 101) showing the low, middle, and top decile for 1.6 km run (1.0 mile) run time for 11,423 girls and boys ages 9-17.

1.6 km run (1.0 mile) run time for 11,423 girls and boys ages 9-17

Age	Male			Female			Male-Female % Difference		
	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile
9	684	522	423	769.0	609.0	499.0	11.1%	14.3%	15.2%
10	666	511	420	759.0	600.0	494.0	12.3%	14.8%	15.0%
11	646	500	416	741.0	586.0	483.0	12.8%	14.7%	13.9%
12	621	485	408	726.0	575.0	474.0	14.5%	15.7%	13.9%
13	587	465	395	716.0	569.0	469.0	18.0%	18.3%	15.8%
14	556	446	382	711.0	567.0	468.0	21.8%	21.3%	18.4%
15	531	432	373	710.0	570.0	469.0	25.2%	24.2%	20.5%
16	514	423	366	710.0	573.0	471.0	27.6%	26.2%	22.3%
17	500	417	362	708.0	575.0	471.0	29.4%	27.5%	23.1%

84. Tomkinson et al. (2018) performed a similarly extensive analysis of literally millions of measurements of a variety of strength and agility metrics from the “Eurofit” test battery on children from 30 European countries. They provide detailed results for each metric, broken out by decile. Sampling the low, middle, and top decile, 9-year-old boys performed better than 9-year-old girls by between 6.5%

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

and 9.7% in the standing broad jump; from 11.4% to 16.1% better in handgrip; and from 45.5% to 49.7% better in the “bent-arm hang.” (Tomkinson 2018.)

85. The Bent Arm Hang test is a measure of upper body muscular strength and endurance used in the Eurofit Physical Fitness Test Battery. To perform the Bent Arm Hang, the child is assisted into position with the body lifted to a height so that the chin is level with the horizontal bar (like a pull up bar). The bar is grasped with the palms facing away from body and the hands shoulder width apart. The timing starts when the child is released. The child then attempts to hold this position for as long as possible. Timing stops when the child's chin falls below the level of the bar, or the head is tilted backward to enable the chin to stay level with the bar.

86. Using data from Tomkinson (2018; table 7 at 1452), the following table sampling the low, middle, and top decile for bent arm hang for 9- to 17-year-old children can be constructed:

Bent Arm Hang time (in seconds) for children ages 9 - 17 years

Age	Male			Female			Male-Female % Difference		
	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile
9	2.13	7.48	25.36	1.43	5.14	16.94	48.95%	45.53%	49.70%
10	2.25	7.92	26.62	1.42	5.15	17.06	58.45%	53.79%	56.04%
11	2.35	8.32	27.73	1.42	5.16	17.18	65.49%	61.24%	61.41%
12	2.48	8.79	28.99	1.41	5.17	17.22	75.89%	70.02%	68.35%
13	2.77	9.81	31.57	1.41	5.18	17.33	96.45%	89.38%	82.17%
14	3.67	12.70	38.39	1.40	5.23	17.83	162.14%	142.83%	115.31%
15	5.40	17.43	47.44	1.38	5.35	18.80	291.30%	225.79%	152.34%
16	7.39	21.75	53.13	1.38	5.63	20.57	435.51%	286.32%	158.29%
17	9.03	24.46	54.66	1.43	6.16	23.61	531.47%	297.08%	131.51%

87. Evaluating these data, a 9-year-old boy in the 50th percentile (that is to say a 9-year-old boy of average upper body muscular strength and endurance) will perform better in the bent arm hang test than 9 through 17-year-old girls in the 50th percentile. Similarly, a 9-year-old boy in the 90th percentile will perform better in the bent arm hang test than 9 through 17-year-old girls in the 90th percentile.

88. Using data from Tomkinson et al. (2017; table 1 at 1549), the following table sampling the low, middle, and top decile for running speed in the last stage of the 20 m shuttle run for 9- to 17-year-old children can be constructed.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

20 m shuttle Running speed (km/h at the last completed stage)

Age	Male			Female			Male-Female % Difference		
	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile
9	8.94	10.03	11.13	8.82	9.72	10.61	1.36%	3.19%	4.90%
10	8.95	10.13	11.31	8.76	9.75	10.74	2.17%	3.90%	5.31%
11	8.97	10.25	11.53	8.72	9.78	10.85	2.87%	4.81%	6.27%
12	9.05	10.47	11.89	8.69	9.83	10.95	4.14%	6.51%	8.58%
13	9.18	10.73	12.29	8.69	9.86	11.03	5.64%	8.82%	11.42%
14	9.32	10.96	12.61	8.70	9.89	11.07	7.13%	10.82%	13.91%
15	9.42	11.13	12.84	8.70	9.91	11.11	8.28%	12.31%	15.57%
16	9.51	11.27	13.03	8.71	9.93	11.14	9.18%	13.49%	16.97%
17	9.60	11.41	13.23	8.72	9.96	11.09	10.09%	14.56%	19.30%

89. Evaluating these data, a 9-year-old boy in the 50th percentile (that is to say a 9-year-old boy of average running speed) will run faster in the final stage of the 20 m shuttle run than 9 through 17-year-old girls in the 50th percentile. Similarly, a 9-year-old boy in the 90th percentile will run faster in the final stage of the 20-m shuttle run than 9 through 15, and 17-year-old girls in the 90th percentile and will be 0.01 km/h (0.01%) slower than 16-year-old girls in the 90th percentile.

90. Just using these two examples for bent arm hang and 20-m shuttle running speed (Tomkinson 2107, Tomkinson 2018) based on large sample sizes (thus having tremendous statistical power) it becomes apparent that a 9-year-old boy will be very likely to outperform similarly trained girls of his own age and older in athletic events involving upper body muscle strength and/or running speed.

91. Another report published in 2014 analyzed physical fitness measurements of 10,302 children aged 6 -10.9 years of age, from the European countries of Sweden, Germany, Hungary, Italy, Cyprus, Spain, Belgium, and Estonia. (De Miguel-Etayo et al. 2014.) The authors observed "... that boys performed better than girls in speed, lower- and upper-limb strength and cardiorespiratory fitness." (57) The data showed that for children of comparable fitness (i.e. 99th percentile boys vs. 99th percentile girls, 50th percentile boys vs. 50th percentile girls, etc.) the boys outperform the girls at every age in measurements of handgrip strength, standing long jump, 20-m shuttle run, and predicted VO₂max (pages 63 and 64, respectively). For clarification, VO₂max is the maximal oxygen consumption, which correlates to 30-40% of success in endurance sports.

92. The standing long jump, also called the Broad Jump, is a common and easy to administer test of explosive leg power used in the Eurofit Physical Fitness Test Battery and in the NFL Combine. In the standing long jump, the participant stands behind a line marked on the ground with feet slightly apart. A two-foot take-

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

off and landing is used, with swinging of the arms and bending of the knees to provide forward drive. The participant attempts to jump as far as possible, landing on both feet without falling backwards. The measurement is taken from takeoff line to the nearest point of contact on the landing (back of the heels) with the best of three attempts being scored.

93. Using data from De Miguel-Etayo et al. (2014, table 3 at 61), which analyzed physical fitness measurements of 10,302 children aged 6 -10.9 years of age, from the European countries of Sweden, Germany, Hungary, Italy, Cyprus, Spain, Belgium, and Estonia, the following table sampling the low, middle, and top decile for standing long jump for 6- to 9-year-old children can be constructed:

Standing Broad Jump (cm) for children ages 6-9 years

Age	Male			Female			Male-Female % Difference		
	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile
6-<6.5	77.3	103.0	125.3	69.1	93.8	116.7	11.9%	9.8%	7.4%
6.5-<7	82.1	108.0	130.7	73.6	98.7	121.9	11.5%	9.4%	7.2%
7-<7.5	86.8	113.1	136.2	78.2	103.5	127.0	11.0%	9.3%	7.2%
7.5-<8	91.7	118.2	141.6	82.8	108.3	132.1	10.7%	9.1%	7.2%
8-<8.5	96.5	123.3	146.9	87.5	113.1	137.1	10.3%	9.0%	7.1%
8.5-<9	101.5	128.3	152.2	92.3	118.0	142.1	10.0%	8.7%	7.1%

94. Another study of Eurofit results for over 400,000 Greek children reported similar results. “[C]ompared with 6-year-old females, 6-year-old males completed 16.6% more shuttle runs in a given time and could jump 9.7% further from a standing position.” (Hilton 2021 at 201, summarizing findings of Tambalis et al. 2016.)

95. Silverman (2011) gathered hand grip data, broken out by age and sex, from a number of studies. Looking only at the nine direct comparisons within individual studies tabulated by Silverman for children aged 7 or younger, in eight of these the boys had strength advantages of between 13 and 28 percent, with the remaining outlier recording only a 4% advantage for 7-year-old boys. (Silverman 2011 Table 1.)

96. To help illustrate the importance of one specific measure of physical fitness in athletic performance, Pocek (2021) stated that to be successful, volleyball “players should distinguish themselves, besides in skill level, in terms of above-average body height, upper and lower muscular power, speed, and agility. Vertical jump is a fundamental part of the spike, block, and serve.” (8377) Pocek further stated that “relative vertical jumping ability is of great importance in volleyball regardless of the players’ position, while absolute vertical jump values can differentiate players not only in terms of player position and performance level but in their career trajectories.” (8382)

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

97. Using data from Ramírez-Vélez (2017; table 2 at 994) which analyzed vertical jump measurements of 7,614 healthy Colombian schoolchildren aged 9 -17.9 years of age the following table sampling the low, middle, and top decile for vertical jump can be constructed:

Vertical Jump Height (cm) for children ages 9 - 17 years

Age	Male			Female			Male-Female % Difference		
	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile
9	18.0	24.0	29.5	16.0	22.3	29.0	12.5%	7.6%	1.7%
10	19.5	25.0	32.0	18.0	24.0	29.5	8.3%	4.2%	8.5%
11	21.0	27.0	32.5	19.5	25.0	31.0	7.7%	8.0%	4.8%
12	22.0	27.5	34.5	20.0	25.5	31.5	10.0%	7.8%	9.5%
13	23.0	30.5	39.0	19.0	25.5	32.0	21.1%	19.6%	21.9%
14	23.5	32.0	41.5	20.0	25.5	32.5	17.5%	25.5%	27.7%
15	26.0	35.5	43.0	20.2	26.0	32.5	28.7%	36.5%	32.3%
16	28.0	36.5	45.1	20.5	26.5	33.0	36.6%	37.7%	36.7%
17	28.0	38.0	47.0	21.5	27.0	35.0	30.2%	40.7%	34.3%

98. Similarly, using data from Taylor (2010; table 2, at 869) which analyzed vertical jump measurements of 1,845 children aged 10 -15 years in primary and secondary schools in the East of England, the following table sampling the low, middle, and top decile for vertical jump can be constructed:

Vertical Jump Height (cm) for children 10 -15 years

Age	Male			Female			Male-Female % Difference		
	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile
10	16.00	21.00	29.00	15.00	22.00	27.00	6.7%	-4.5%	7.4%
11	20.00	27.00	34.00	19.00	25.00	32.00	5.3%	8.0%	6.3%
12	23.00	30.00	37.00	21.00	27.00	33.00	9.5%	11.1%	12.1%
13	23.00	32.00	40.00	21.00	26.00	34.00	9.5%	23.1%	17.6%
14	26.00	36.00	44.00	21.00	28.00	34.00	23.8%	28.6%	29.4%
15	29.00	37.00	44.00	21.00	28.00	39.00	38.1%	32.1%	12.8%

99. As can be seen from the data from Ramírez-Vélez (2017) and Taylor (2010), males consistently outperform females of the same age and percentile in vertical jump height. Both sets of data show that an 11-year-old boy in the 90th percentile for vertical jump height will outperform girls in the 90th percentile at ages 11 and 12, and will be equal to girls at ages 13, 14, and possibly 15. These data indicate that an 11-year-old would be likely to have an advantage over girls of the same age and older in sports such as volleyball where “absolute vertical jump

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

values can differentiate players not only in terms of player position and performance level but in their career trajectories.” (Pocek 2021 at 8382.)

100. Boys also enjoy an advantage in throwing well before puberty. “Boys exceed girls in throwing velocity by 1.5 standard deviation units as early as 4 to 7 years of age. . . The boys exceed the girls [in throwing distance] by 1.5 standard deviation units as early as 2 to 4 years of age.” (Thomas 1985 at 266.) This means that the average 4- to 7-year-old boy can out-throw approximately 87% of all girls of his age.

101. Record data from USA Track & Field indicate that boys outperform girls in track events even in the youngest age group for whom records are kept (age 8 and under).⁸

**American Youth Outdoor Track & Field Record times in
age groups 8 and under (time in seconds)**

Event	Boys	Girls	Difference
100M	13.65	13.78	0.95%
200M	27.32	28.21	3.26%
400M	62.48	66.10	5.79%
800M	148.59	158.11	6.41%
1500M	308.52	314.72	2.01%
Mean			3.68%

102. Looking at the best times within a single year shows a similar pattern of consistent advantage for even young boys. I consider the 2018 USATF Region 8 Junior Olympic Championships for the youngest age group (8 and under).⁹

2018 USATF Region 8 Junior Olympic Championships for the 8 and under age group

Event	Boys	Girls	Difference
100M	15.11	15.64	3.51%
200M	30.79	33.58	9.06%
400M	71.12	77.32	8.72%
800M	174.28	180.48	3.56%
1500M	351.43	382.47	8.83%
Mean			6.74%

⁸<http://legacy.usatf.org/statistics/records/view.asp?division=american&location=outdoor%20track%20%26%20field&age=youth&sport=TF>

⁹ <https://www.athletic.net/TrackAndField/meet/384619/results/m/1/100m>

⁹ <https://www.athletic.net/CrossCountry/Division/List.aspx?DivID=62211>

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

103. Using Athletic.net⁹, for 2021 Cross Country and Track & Field data for boys and girls in the 7-8, 9-10, and 11-12 year old age group club reports, and for 5th, 6th, and 7th grade for the whole United States I have compiled the tables for 3000 m events, and for the 100-m, 200-m, 400-m, 800-m, 1600-m, 3000-m, long jump, and high jump Track and Field data to illustrate the differences in individual athletic performance between boys and girls, all of which appear in the Appendix. The pattern of males outperforming females was consistent across events, with rare anomalies, only varying in the magnitude of difference between males and females.

104. Similarly, using Athletic.net, for 2021 Track & Field data for boys and girls in the 6th grade for the state of West Virginia, I have compiled tables, which appear in the appendix, comparing the performance of boys and girls for the 100-m, 200-m, 400-m, 800-m, 1600-m, and 3200-m running events in which the 1st place boy was consistently faster than the 1st place girl, and the average performance of the top 10 boys was consistently faster than the average performance for the top 10 girls. Based on the finishing times for the 1st place boy and girl in the 6th grade in West Virginia 1600-m race, and extrapolating the running time to a running pace, the 1st place boy would be expected to finish 273 m in front of the 1st place girl, which is 2/3 of a lap on a standard 400-m track, or almost the length of 3 football fields. In comparison, the 1st place boy would finish 66 m in front of the 2nd place boy, and the 1st place girl would finish 20 m in front of the 2nd place girl.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Top 10 West Virginia boys and girls 6th grade outdoor track for 2021 (time in seconds)

100 m			200 m			400 m		
	Boys	Girls		Boys	Girls		Boys	Girls
1	13.18	14.00	Difference	26.97	29.28	Difference	60.04	65.50
2	13.94	14.19	between #1	29.38	30.05	between #1	60.48	67.51
3	14.07	14.47	boy and # 1	30.09	30.34	boy and # 1	66.26	68.60
4	14.44	14.86	girl	30.10	30.73	girl	67.12	70.43
5	14.46	14.92	5.9%	30.24	31.00	7.9%	68.28	71.09
6	14.53	15.04		30.38	31.04		68.36	71.38
7	14.75	15.04	Average	30.54	31.10	Average	69.65	73.61
8	14.78	15.20	difference	30.69	31.10	difference	69.70	73.87
			boys vs			boys vs		
9	14.84	15.25	girls	30.74	31.35	girls	69.76	74.07
10	14.94	15.28	2.9%	30.99	31.64	2.4%	70.63	74.21
800 m			1600 m			3200 m		
	Boys	Girls		Boys	Girls		Boys	Girls
1	147.2	164.5	Difference	305.5	357.8	Difference	678.4	776.6
2	147.9	166.1	between #1	318.1	361.6	between #1	750.0	809.8
3	152.1	167.2	boy and # 1	322.0	379.8	boy and # 1	763.3	811.0
4	153.2	170.2	girl	336.0	385.2	girl	766.3	843.0
5	155.3	171.0	10.6%	342.2	390.2	14.6%	771.7	850.6
6	159.5	171.5		348.0	392.0		782.8	852.1
7	159.9	174.8	Average	356.6	393.3	Average	794.1	858.0
8	167.8	174.9	difference	357.5	395.7	difference	803.0	862.8
			boys vs			boys vs		
9	169.2	175.9	girls	362.4	398.1	girls	812.1	869.9
10	172.6	177.6	7.5%	366.0	403.2	11.5%	814.3	883.3

105. As serious runners will recognize, differences of 3%, 5%, or 8% are not easily overcome. During track competition the difference between first and second place, or second and third place, or third and fourth place (and so on) is often 0.5 - 0.7%, with some contests being determined by as little as 0.01%.

106. I performed an analysis of running events (consisting of the 100-m, 200-m, 400-m, 800-m, 1500-m, 5000-m, and 10,000-m) in the Division 1, Division 2, and Division 3 NCAA Outdoor championships for the years of 2010-2019: the mean difference between 1st and 2nd place was 0.48% for men and 0.86% for women. The mean difference between 2nd and 3rd place was 0.46% for men and 0.57% for women. The mean difference between 3rd place and 4th place was 0.31% for men and 0.44% for women. The mean difference between 1st place and 8th place (the last place to earn the title of All American) was 2.65% for men and 3.77% for women. (Brown et al. Unpublished observations, to be presented at the 2022 Annual Meeting of the American College of Sports Medicine.)

107. A common response to empirical data showing pre-pubertal performance advantages in boys is the argument that the performance of boys may

represent a social–cultural bias for boys to be more physically active, rather than representing inherent sex-based differences in pre-pubertal physical fitness. However, the younger the age at which such differences are observed, and the more egalitarian the culture within which they are observed, the less plausible this hypothesis becomes. Eiberg et al. (2005) measured body composition, VO₂max, and physical activity in 366 Danish boys and 332 Danish girls between the ages of 6 and 7 years old. Their observations indicated that VO₂max was 11% higher in boys than girls. When expressed relative to body mass the boys' VO₂max was still 8% higher than the girls. The authors stated that “...no differences in haemoglobin or sex hormones¹⁰ have been reported in this age group,” yet “... when children with the same VO₂max were compared, boys were still more active, and in boys and girls with the same P[hysical] A[ctivity] level, boys were fitter.” (728). These data indicate that in pre-pubertal children, in a very egalitarian culture regarding gender roles and gender norms, boys still have a measurable advantage in regards to aerobic fitness when known physiological and physical activity differences are accounted for.

108. And, as I have mentioned above, even by the age of 4 or 5, in a ruler-drop test, boys exhibit 4% to 6% faster reaction times than girls. (Latorre-Roman 2018.)

109. When looking at the data on testosterone concentrations previously presented, along with the data on physical fitness and athletic performance presented, boys have advantages in athletic performance and physical fitness before there are marked differences in testosterone concentrations between boys and girls.

110. For the most part, the data I review above relate to pre-pubertal children. Today, we also face the question of inclusion in female athletics of males who have undergone “puberty suppression.” The UK Sport Councils Literature Review notes that, “In the UK, so-called ‘puberty blockers’ are generally not used until Tanner maturation stage 2-3 (i.e. after puberty has progressed into early sexual maturation).” (9.) While it is outside my expertise, my understanding is that current practice with regard to administration of puberty blockers is similar in the United States. Tanner stages 2 and 3 generally encompass an age range from 10 to 14 years old, with significant differences between individuals. Like the authors of the UK Sports Council Literature Review, I am “not aware of research” directly addressing the implications for athletic capability of the use of puberty blockers. (UK Sport Councils Literature Review at 9.) As Handelsman documents, the male advantage begins to increase rapidly—along with testosterone levels—at about age 11, or “very closely aligned to the timing of the onset of male puberty.” (Handelsman 2017.) It seems likely that males who have undergone puberty suppression will

¹⁰ This term would include testosterone and estrogens.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

have physiological and performance advantages over females somewhere between those possessed by pre-pubertal boys, and those who have gone through full male puberty, with the degree of advantage in individual cases depending on that individual's development and the timing of the start of puberty blockade.

111. Tack et al. (2018) observed that in 21 transgender-identifying biological males, administration of antiandrogens for 5-31 months (commencing at 16.3 ± 1.21 years of age), resulted in nearly, but not completely, halting of normal age-related *increases* in muscle strength. Importantly, muscle strength did not decrease after administration of antiandrogens. Rather, despite antiandrogens, these individuals retained higher muscle mass, lower percent body fat, higher body mass, higher body height, and higher grip strength than comparable girls of the same age. (Supplemental tables).

112. Klaver et al. (2018 at 256) demonstrated that the use of puberty blockers did not eliminate the differences in lean body mass between biological male and female teenagers. Subsequent use of puberty blockers combined with cross-sex hormone use (in the same subjects) still did not eliminate the differences in lean body mass between biological male and female teenagers. Furthermore, by 22 years of age, the use of puberty blockers, and then puberty blockers combined with cross sex hormones, and then cross hormone therapy alone for over 8 total years of treatment still had not eliminated the difference in lean body mass between biological males and females.

113. The effects of puberty blockers on growth and development, including muscle mass, fat mass, or other factors that influence athletic performance, have been minimally researched. Indeed, Klaver et al. (2018) is the only published research that I am aware of that has evaluated the use of puberty blockers on body composition. As stated by Roberts and Carswell (2021), "No published studies have fully characterized the impact of [puberty blockers on] final adult height or current height in an actively growing TGD youth." (1680). Likewise, "[n]o published literature provides guidance on how to best predict the final adult height for TGD youth receiving GnRHa and gender-affirming hormonal treatment." (1681). Thus, the effect of prescribing puberty blockers to a male child before the onset of puberty on the physical components of athletic performance is largely unknown. There is not any scientific evidence that such treatment eliminates the pre-existing performance advantages that prepubertal males have over prepubertal females.

B. The rapid increase in testosterone across male puberty drives characteristic male physiological changes and the increasing performance advantages.


114. While boys exhibit some performance advantage even before puberty, it is both true and well known to common experience that the male advantage

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

increases rapidly, and becomes much larger, as boys undergo puberty and become men. Empirically, this can be seen by contrasting the modest advantages reviewed immediately above against the large performance advantages enjoyed by men that I have detailed in Section II.

115. Multiple studies (along with common observation) document that the male performance advantage begins to increase during the early years of puberty, and then increases rapidly across the middle years of puberty (about ages 12-16). (Tønnessen 2015; Handelsman 2018 at 812-813.) Since it is well known that testosterone levels increase by more than an order of magnitude in boys across puberty, it is unsurprising that Handelsman finds that these increases in male performance advantage correlate to increasing testosterone levels, as presented in his chart reproduced below. (Handelsman 2018 at 812-13.)



116. Handelsman further finds that certain characteristic male changes including boys' increase in muscle mass do not begin at all until "circulating testosterone concentrations rise into the range of males at mid-puberty, which are higher than in women at any age." (Handelsman 2018 at 810.)

117. Knox et al. (2019) agree that "[i]t is well recognised that testosterone contributes to physiological factors including body composition, skeletal structure, and the cardiovascular and respiratory systems across the life span, with significant influence during the pubertal period. These physiological factors underpin strength, speed, and recovery with all three elements required to be competitive in almost all sports." (Knox 2019 at 397.) "High testosterone levels and prior male physiology provide an all-purpose benefit, and a substantial advantage. As the IAAF says, "To the best of our knowledge, there is no other genetic or biological trait encountered in female athletics that confers such a huge performance advantage." (Knox 2019 at 399.)

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

118. However, the undisputed fact that high (that is, normal male) levels of testosterone drive the characteristically male physiological changes that occur across male puberty does not at all imply that artificially *depressing* testosterone levels after those changes occur will reverse all or most of those changes so as to eliminate the male athletic advantage. This is an empirical question. As it turns out, the answer is that while some normal male characteristics can be changed by means of testosterone suppression, others cannot be, and all the reliable evidence indicates that males retain large athletic advantages even after long-term testosterone suppression.

V. The available evidence shows that suppression of testosterone in a male after puberty has occurred does not substantially eliminate the male athletic advantage.

119. The 2011 “NCAA Policy on Transgender Student-Athlete Participation” requires only that males who identify as transgender be on unspecified and unquantified “testosterone suppression treatment” for “one calendar year” prior to competing in women’s events. In supposed justification of this policy, the NCAA’s Office of Inclusion asserts that, “It is also important to know that any strength and endurance advantages a transgender woman arguably may have as a result of her prior testosterone levels dissipate after about one year of estrogen or testosterone-suppression therapy.” (NCAA 2011 at 8.)

120. Similarly, writing in 2018, Handelsman et al. could speculate that even though some male advantages established during puberty are “fixed and irreversible (bone size),” “[t]he limited available prospective evidence . . . suggests that the advantageous increases in muscle and hemoglobin due to male circulating testosterone concentrations are induced or reversed during the first 12 months.” (Handelsman 2018 at 824.)

121. But these assertions or hypotheses of the NCAA and Handelsman are now strongly contradicted by the available science. In this section, I examine what is known about whether suppression of testosterone in males can eliminate the male physiological and performance advantages over females.

A. Empirical studies find that males retain a strong performance advantage even after lengthy testosterone suppression.

122. As my review in Section II indicates, a very large body of literature documents the large performance advantage enjoyed by males across a wide range of athletics. To date, only a limited number of studies have directly measured the effect of testosterone suppression and the administration of female hormones on the athletic performance of males. These studies report that testosterone suppression for a full year (and in some cases much longer) does not come close to eliminating

male advantage in strength (hand grip, leg strength, and arm strength) or running speed.

Hand Grip Strength

123. As I have noted, hand grip strength is a well-accepted proxy for general strength. Multiple separate studies, from separate groups, report that males retain a large advantage in hand strength even after testosterone suppression to female levels.

124. In a longitudinal study, Van Caenegem et al. reported that males who underwent standard testosterone suppression protocols lost only 7% hand strength after 12 months of treatment, and only a cumulative 9% after two years. (Van Caenegem 2015 at 42.) As I note above, on average men exhibit in the neighborhood of 60% greater hand grip strength than women, so these small decreases do not remotely eliminate that advantage. Van Caenegem et al. document that their sample of males who elected testosterone suppression began with less strength than a control male population. Nevertheless, after one year of suppression, their study population still had hand grip only 21% less than the control male population, and thus still far higher than a female population. (Van Caenegem 2015 at 42.)

125. Scharff et al. (2019) measured grip strength in a large cohort of male-to-female subjects from before the start of hormone therapy through one year of hormone therapy. The hormone therapy included suppression of testosterone to less than 2 nmol/L “in the majority of the transwomen,” (1024), as well as administration of estradiol (1021). These researchers observed a small decrease in grip strength in these subjects over that time (Fig. 2), but mean grip strength of this group remained far higher than mean grip strength of females—specifically, “After 12 months, the median grip strength of transwomen [male-to-female subjects] still falls in the 95th percentile for age-matched females.” (1026).

126. Still a third longitudinal study, looking at teen males undergoing testosterone suppression, “noted no change in grip strength after hormonal treatment (average duration 11 months) of 21 transgender girls.” (Hilton 2021 at 207, summarizing Tack 2018.)

127. In a fourth study, Lapauw et al. (2008) looked at the extreme case of testosterone suppression by studying a population of 23 biologically male individuals who had undergone at least two years of testosterone suppression, followed by sex reassignment surgery that included “orchidectomy” (that is, surgical castration), and then at least an additional three years before the study date. Comparing this group against a control of age- and height-matched healthy males, the researchers found that the individuals who had gone through testosterone suppression and then surgical castration had an average hand grip (41 kg) that was

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

24% weaker than the control group of healthy males. But this remains at least 25% *higher* than the average hand-grip strength of biological females as measured by Bohannon et al. (2019).

128. Summarizing these and a few other studies measuring strength loss (in most cases based on hand grip) following testosterone suppression, Harper et al. (2021) conclude that “strength loss with 12 months of [testosterone suppression] . . . ranged from non-significant to 7%. . . . [T]he small decrease in strength in transwomen after 12-36 months of [testosterone suppression] suggests that transwomen likely retain a strength advantage over cisgender women.” (Hilton 2021 at 870.)

Arm Strength

129. Lapauw et al. (2008) found that 3 years after surgical castration, preceded by at least two years of testosterone suppression, biologically male subjects had 33% less bicep strength than healthy male controls. (Lapauw (2008) at 1018.) Given that healthy men exhibit between 89% and 109% greater arm strength than healthy women, this leaves a very large residual arm strength advantage over biological women.

130. Roberts et al. have recently published an interesting longitudinal study, one arm of which considered biological males who began testosterone suppression and cross-sex hormones while serving in the United States Air Force. (Roberts 2020.) One measured performance criterion was pushups per minute, which, while not exclusively, primarily tests arm strength under repetition. *Before* treatment, the biological male study subjects who underwent testosterone suppression could do 45% more pushups per minute than the average for all Air Force women under the age of 30 (47.3 vs. 32.5). *After* between one and two years of testosterone suppression, this group could still do 33% more pushups per minute. (Table 4.) Further, the body weight of the study group did not decline at all after one to two years of testosterone suppression (in fact rose slightly) (Table 3), and was approximately 24 pounds (11.0 kg) higher than the average for Air Force women under the age of 30. (Roberts 2020 at 3.) This means that the individuals who had undergone at least one year of testosterone suppression were not only doing 1/3 more pushups per minute, but were lifting significantly more weight with each pushup.

131. After two years of testosterone suppression, the study sample in Roberts et al. was only able to do 6% more pushups per minute than the Air Force female average. But their weight remained unchanged from their pre-treatment starting point, and thus about 24 pounds higher than the Air Force female average. As Roberts et al. explain, “as a group, transwomen weigh more than CW [cis-women]. Thus, transwomen will have a higher power output than CW when

performing an equivalent number of push-ups. Therefore, our study may underestimate the advantage in strength that transwomen have over CW.” (Roberts 2020 at 4.)

Leg Strength

132. Wiik et al. (2020), in a longitudinal study that tracked 11 males from the start of testosterone suppression through 12 months after treatment initiation, found that isometric strength levels measured at the knee “were maintained over the [study period].”¹¹ (808) “At T12 [the conclusion of the one-year study], the absolute levels of strength and muscle volume were greater in [male-to-female subjects] than in . . . CW [women who had not undergone any hormonal therapy].” (Wiik 2020 at 808.) In fact, Wiik et al. reported that “muscle strength after 12 months of testosterone suppression was comparable to baseline strength. As a result, transgender women remained about 50% stronger than . . . a reference group of females.” (Hilton 2021 at 207, summarizing Wiik 2020.)

133. Lapauw et al. (2008) found that 3 years after surgical castration, preceded by at least two years of testosterone suppression, subjects had peak knee torque only 25% lower than healthy male controls. (Lapauw 2008 at 1018.) Again, given that healthy males exhibit 54% greater maximum knee torque than healthy females, this leaves these individuals with a large average strength advantage over females even years after sex reassignment surgery.

Running speed

134. The most striking finding of the recent Roberts et al. study concerned running speed over a 1.5 mile distance—a distance that tests midrange endurance. Before suppression, the MtF study group ran 21% faster than the Air Force female average. After at least 2 year of testosterone suppression, these subjects still ran 12% faster than the Air Force female average. (Roberts 2020 Table 4.)

135. The specific experience of the well-known case of NCAA athlete Cece Telfer is consistent with the more statistically meaningful results of Roberts et al., further illustrating that male-to-female transgender treatment does not negate the inherent athletic performance advantages of a post-pubertal male. In 2016 and 2017 Cece Telfer competed as Craig Telfer on the Franklin Pierce University men’s track team, being ranked 200th and 390th (respectively) against other NCAA Division 2 men. “Craig” Telfer did not qualify for the National Championships in any events. Telfer did not compete in the 2018 season while undergoing testosterone

¹¹ Isometric strength measures muscular force production for a given amount of time at a specific joint angle but with no joint movement.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

suppression (per NCAA policy). In 2019 Cece Telfer competed on the Franklin Pierce University *women's* team, qualified for the NCAA Division 2 Track and Field National Championships, and placed 1st in the women's 400 meter hurdles and placed third in the women's 100 meter hurdles. (For examples of the media coverage of this please see <https://www.washingtontimes.com/news/2019/jun/3/cece-telfer-franklin-pierce-transgenderhurdler-wi/> last accessed May 29, 2020. <https://www.newshub.co.nz/home/sport/2019/06/athletics-transgender-woman-cece-telfer-whopreviously-competed-as-a-man-wins-ncaa-track-championship.html> (last accessed May 29, 2020).)

136. The table below shows the best collegiate performance times from the combined 2015 and 2016 seasons for Cece Telfer when competing as a man in men's events, and the best collegiate performance times from the 2019 season when competing as a woman in women's events. Comparing the times for the running events (in which male and female athletes run the same distance) there is no statistical difference between Telfer's "before and after" times. Calculating the difference in time between the male and female times, Telfer performed an average of 0.22% *faster* as a female. (Comparing the performance for the hurdle events (marked with H) is of questionable validity due to differences between men's and women's events in hurdle heights and spacing, and distance for the 110m vs. 100 m.) While this is simply one example, and does not represent a controlled experimental analysis, this information provides some evidence that male-to-female transgender treatment does not negate the inherent athletic performance advantages of a postpubertal male. (These times were obtained from https://www.tfrrs.org/athletes/6994616/Franklin_Pierce/CeCe_Telfer.html and <https://www.tfrrs.org/athletes/5108308.html>, last accessed May 29, 2020).

As Craig Telfer (male athlete)		As Cece Telfer (female athlete)	
Event	Time (seconds)	Event	Time (seconds)
55	7.01	55	7.02
60	7.67	60	7.63
100	12.17	100	12.24
200	24.03	200	24.30
400	55.77	400	54.41
55 H †	7.98	55 H †	7.91
60 H †	8.52	60 H †	8.33
110 H †	15.17	100 H †	13.41*
400 H ‡	57.34	400 H ‡	57.53**

* women's 3rd place, NCAA Division 2 National Championships

** women's 1st place, NCAA Division 2 National Championships

† men's hurdle height is 42 inches with differences in hurdle spacing between men and women

‡ men's hurdle height is 36 inches, women's height is 30 inches with the same spacing between hurdles

137. Similarly, University of Pennsylvania swimmer Lia Thomas began competing in the women's division in the fall of 2021, after previously competing for U. Penn. in the men's division. Thomas has promptly set school, pool, and/or league women's records in 200 yard freestyle, 500 yard freestyle, and 1650 yard freestyle competitions, beating the nearest female in the 1650 yard by an unheard-of 38 seconds.

138. In a pre-peer review article, Senefeld, Coleman, Hunter, and Joyner (doi: <https://doi.org/10.1101/2021.12.28.21268483>, accessed January 12, 2022) "compared the gender-related differences in performance of a transgender swimmer who competed in both the male and female NCAA (collegiate) categories to the sex-related differences in performance of world and national class swimmers" and observed that this athlete [presumably Lia Thomas based on performance times and the timing of this article] was unranked in 2018-2019 in the 100-yard, ranked 551st in the 200-yard, 65th in the 500-yard 32nd in the 1650-yards men's freestyle. After following the NCAA protocol for testosterone suppression and competing as a woman in 2021-2022, this swimmer was ranked 94th in the 100-yard, 1st in the 200-yard, 1st in the 500-yard, and 6th in the 1650-yard women's freestyle. The performance times swimming as a female, when compared to swimming as a male, were 4.6% slower in the 100-yard, 2.6% slower in the 200-yard, 5.6% slower in the 500-yard, and 6.8% slower in the 1650-yard events than when swimming as a male. *It is important to note that these are mid-season race times and do not represent season best performance times or in a championship event where athletes often set their personal record times.* The authors concluded "...that for middle distance events (100, 200 and 400m or their imperial equivalents) lasting between about one and five minutes, the decrements in performance of the transgender woman swimmer are less than expected on the basis of a comparison of a large cohort of world and national class performances by female and male swimmers" and "it is possible that the relative improvements in this swimmer's rankings in the women's category relative to the men's category are due to legacy effects of testosterone on a number of physiological factors that can influence athletic performance."

139. Harper (2015) has often been cited as "proving" that testosterone suppression eliminates male advantage. And indeed, hedged with many disclaimers, the author in that article does more or less make that claim with respect to "distance races," while emphasizing that "the author makes no claims as to the equality of performances, pre and post gender transition, in any other sport." (Harper 2015 at 8.) However, Harper (2015) is in effect a collection of unverified anecdotes, not science. It is built around self-reported race times from just eight self-selected transgender runners, recruited "mostly" online. How and on what websites the subjects were recruited is not disclosed, nor is anything said about how those not recruited online were recruited. Thus, there is no information to tell us whether these eight runners could in any way be representative, and the

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

recruitment pools and methodology, which could bear on ideological bias in their self-reports, is not disclosed.

140. Further, the self-reported race times relied on by Harper (2015) *span 29 years*. It is well known that self-reported data, particularly concerning emotionally or ideologically fraught topics, is unreliable, and likewise that memory of distant events is unreliable. Whether the subjects were responding from memory or from written records, and if so what records, is not disclosed, and does not appear to be known to the author. For six of the subjects, the author claims to have been able to verify “approximately half” of the self-reported times. Which scores these are is not disclosed. The other two subjects responded only anonymously, so nothing about their claims could be or was verified. In short, neither the author nor the reader knows whether the supposed “facts” on which the paper’s analysis is based are true.

141. Even if we could accept them at face value, the data are largely meaningless. Only two of the eight study subjects reported (undefined) “stable training patterns,” and even with consistent training, athletic performance generally declines with age. As a result, when the few data points span 29 years, it is not possible to attribute declines in performance to asserted testosterone suppression. Further, distance running is usually not on a track, and race times vary significantly depending on the course and the weather. Only one reporting subject who claimed a “stable training pattern” reported “before and after” times on the same course within three years’ time,” which the author acknowledges would “represent the best comparison points.”

142. Harper (2015) to some extent acknowledges its profound methodological flaws, but seeks to excuse them by the difficulty of breaking new ground. The author states that, “The first problem is how to formulate a study to create a meaningful measurement of athletic performance, both before and after testosterone suppression. No methodology has been previously devised to make meaningful measurements.” (2) This statement was not accurate at the time of publication, as there are innumerable publications with validated methodology for comparing physical fitness and/or athletic performance between people of different ages, sexes, and before and after medical treatment, any of which could easily have been used with minimal or no adaptation for the purposes of this study. Indeed, well before the publication of Harper (2015), several authors that I have cited in this review had performed and published disciplined and methodologically reliable studies of physical performance and physiological attributes “before and after” testosterone suppression.

143. More recently, and to her credit, Harper has acknowledged the finding of Roberts (2020) regarding the durable male advantage in running speed in the 1.5 mile distance, even after two years of testosterone suppression. She joins with co-

authors in acknowledging that this study of individuals who (due to Air Force physical fitness requirements) “could at least be considered exercise trained,” agrees that Roberts’ data shows that “transwomen ran significantly faster during the 1.5 mile fitness test than ciswomen,” and declares that this result is “consistent with the findings of the current review in untrained transgender individuals” that even 30 months of testosterone suppression does not eliminate all male advantages “associated with muscle endurance and performance.” (Harper 2021 at 8.) The Harper (2021) authors conclude overall “that strength may be well preserved in transwomen during the first 3 years of hormone therapy,” and that [w]hether transgender and cisgender women can engage in meaningful sport [in competition with each other], even after [testosterone suppression], is a highly debated question.” (Harper 2021 at 1, 8.)

144. Higerd (2021) “[a]ssess[ed] the probability of a girls’ champion being biologically male” by evaluating 920,11 American high school track and field performances available through the track and field database Athletic.net in five states (CA, FL, MN, NY, WA), over three years (2017 – 2019), in eight events; high jump, long jump, 100M, 200M, 400M, 800M, 1600M, and 3200M and estimated that “there is a simulated 81%-98% probability of transgender dominance occurring in the female track and field event” and further concluded that “in the majority of cases, the entire podium (top of the state) would be MTF [transgender athletes]” (at xii).

B. Testosterone suppression does not reverse important male physiological advantages.

145. We see that, once a male has gone through male puberty, later testosterone suppression (or even castration) leaves large strength and performance advantages over females in place. It is not surprising that this is so. What is now a fairly extensive body of literature has documented that many of the specific male physiological advantages that I reviewed in Section II are not reversed by testosterone suppression after puberty, or are reduced only modestly, leaving a large advantage over female norms still in place.

146. Handelsman has well documented that the large increases in physiological and performance advantages characteristic of men develop in tandem with, and are likely driven by, the rapid and large increases in circulating testosterone levels that males experience across puberty, or generally between the ages of about 12 through 18. (Handelsman 2018.) Some have misinterpreted Handelsman as suggesting that all of those advantages are and remain entirely dependent—on an ongoing basis—on *current* circulating testosterone levels. This is a misreading of Handelsman, who makes no such claim. As the studies reviewed above demonstrate, it is also empirically false with respect to multiple measures of

performance. Indeed, Handelsman himself, referring to the Roberts et al. (2020) study which I describe below, has recently written that “transwomen treated with estrogens after completing male puberty experienced only minimal declines in physical performance over 12 months, substantially surpassing average female performance for up to 8 years.” (Handelsman 2020.)

147. As to individual physiological advantages, the more accurate and more complicated reality is reflected in a statement titled “The Role of Testosterone in Athletic Performance,” published in 2019 by several dozen sports medicine experts and physicians from many top medical schools and hospitals in the U.S. and around the world. (Levine et al. 2019.) This expert group concurs with Handelsman regarding the importance of testosterone to the male advantage, but recognizes that those advantages depend not only on *current* circulating testosterone levels in the individual, but on the “exposure in biological males to much higher levels of testosterone during growth, development, and throughout the athletic career.” (*Emphasis added.*) In other words, both past and current circulating testosterone levels affect physiology and athletic capability.

148. Available research enables us to sort out, in some detail, which specific physiological advantages are immutable once they occur, which can be reversed only in part, and which appear to be highly responsive to later hormonal manipulation. The bottom line is that very few of the male physiological advantages I have reviewed in Section II above are largely reversible by testosterone suppression once an individual has passed through male puberty.

Skeletal Configuration

149. It is obvious that some of the physiological changes that occur during “growth and development” across puberty cannot be reversed. Some of these irreversible physiological changes are quite evident in photographs that have recently appeared in the news of transgender competitors in female events. These include skeletal configuration advantages including:

- Longer and larger bones that give height, weight, and leverage advantages to men;
- More advantageous hip shape and configuration as compared to women.

Cardiovascular Advantages

150. Developmental changes for which there is no apparent means of reversal, and no literature suggesting reversibility, also include multiple

contributors to the male cardiovascular advantage, including diaphragm placement, lung and trachea size, and heart size and therefore pumping capacity.¹²

151. On the other hand, the evidence is mixed as to hemoglobin concentration, which as discussed above is a contributing factor to $\dot{V}O_2$ max. Harper (2021) surveyed the literature and found that “Nine studies reported the levels of Hgb [hemoglobin] or HCT [red blood cell count] in transwomen before and after [testosterone suppression], from a minimum of three to a maximum of 36 months post hormone therapy. Eight of these studies. . . found that hormone therapy led to a significant (4.6%–14.0%) decrease in Hgb/HCT ($p < 0.01$), while one study found no significant difference after 6 months,” but only one of those eight studies returned results at the generally accepted 95% confidence level. (Harper 2021 at 5-6 and Table 5.)

152. I have not found any study of the effect of testosterone suppression on the male advantage in mitochondrial biogenesis.

Muscle mass

153. Multiple studies have found that muscle mass decreases modestly or not at all in response to testosterone suppression. Knox et al. report that “healthy young men did not lose significant muscle mass (or power) when their circulating testosterone levels were reduced to 8.8 nmol/L (lower than the 2015 IOC guideline of 10 nmol/L) for 20 weeks.” (Knox 2019 at 398.) Gooren found that “[i]n spite of muscle surface area reduction induced by androgen deprivation, after 1 year the mean muscle surface area in male-to- female transsexuals remained significantly greater than in untreated female-to-male transsexuals.” (Gooren 2011 at 653.) An earlier study by Gooren found that after one year of testosterone suppression, muscle mass at the thigh was reduced by only about 10%, exhibited “no further reduction after 3 years of hormones,” and “remained significantly greater” than in his sample of untreated women. (Gooren 2004 at 426-427.) Van Caenegem et al. found that muscle cross section in the calf and forearm decreased only trivially (4% and 1% respectively) after two years of testosterone suppression. (Van Caenegem 2015 Table 4.)

154. Taking measurements one month after start of testosterone suppression in male-to-female (non-athlete) subjects, and again 3 and 11 months after start of feminizing hormone replacement therapy in these subjects, Wiik et al.

¹² “[H]ormone therapy will not alter ... lung volume or heart size of the transwoman athlete, especially if [that athlete] transitions postpuberty, so natural advantages including joint articulation, stroke volume and maximal oxygen uptake will be maintained.” (Knox 2019 at 398.)

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

found that total lean tissue (i.e. primarily muscle) did not decrease significantly across the entire period. Indeed, “some of the [subjects] did not lose any muscle mass at all.” (Wiik 2020 at 812.) And even though they observed a small decrease in thigh muscle mass, they found that isometric strength levels measured at the knee “were maintained over the [study period].” (808) “At T12 [the conclusion of the one-year study], the absolute levels of strength and muscle volume were greater in [male-to-female subjects] than in [female-to-male subjects] and CW [women who had not undergone any hormonal therapy].” (808)

155. Hilton & Lundberg summarize an extensive survey of the literature as follows:

“12 longitudinal studies have examined the effects of testosterone suppression on lean body mass or muscle size in transgender women. The collective evidence from these studies suggests that 12 months, which is the most commonly examined intervention period, of testosterone suppression to female typical reference levels results in a modest (approximately– 5%) loss of lean body mass or muscle size. . . .

“Thus, given the large baseline differences in muscle mass between males and females (Table 1; approximately 40%), the reduction achieved by 12 months of testosterone suppression can reasonably be assessed as small relative to the initial superior mass. We, therefore, conclude that the muscle mass advantage males possess over females, and the performance implications thereof, are not removed by the currently studied durations (4 months, 1, 2 and 3 years) of testosterone suppression in transgender women. (Hilton 2021 at 205-207.)

156. When we recall that “women have 50% to 60% of men’s upper arm muscle cross-sectional area and 65% to 70% of men’s thigh muscle cross-sectional area” (Handelsman 2018 at 812), it is clear that Hilton’s conclusion is correct. In other words, biologically male subjects possess substantially larger muscles than biologically female subjects after undergoing a year or even three years of testosterone suppression.

157. I note that outside the context of transgender athletes, the testosterone-driven increase in muscle mass and strength enjoyed by these male-to-female subjects would constitute a disqualifying doping violation under all league anti-doping rules with which I am familiar.

C. Responsible voices internationally are increasingly recognizing that suppression of testosterone in a male after puberty has occurred does not substantially reverse the male athletic advantage.

158. The previous very permissive NCAA policy governing transgender participation in women's collegiate athletics was adopted in 2011, and the previous IOC guidelines were adopted in 2015. At those dates, much of the scientific analysis of the actual impact of testosterone suppression had not yet been performed, much less any wider synthesis of that science. In fact, a series of important peer-reviewed studies and literature reviews have been published only very recently, since I prepared my first paper on this topic, in early 2020.

159. These new scientific publications reflect a remarkably consistent consensus: once an individual has gone through male puberty, testosterone suppression does not substantially eliminate the physiological and performance advantages that that individual enjoys over female competitors.

160. Importantly, I have found no peer-reviewed scientific paper, nor any respected scientific voice, that is now asserting the contrary—that is, that testosterone suppression can eliminate or even largely eliminate the male biological advantage once puberty has occurred.

161. I excerpt the key conclusions from important recent peer-reviewed papers below.

162. Roberts 2020: “In this study, we confirmed that . . . the pretreatment differences between transgender and cis gender women persist beyond the 12-month time requirement currently being proposed for athletic competition by the World Athletics and the IOC.” (6)

163. Wiik 2020: The muscular and strength changes in males undergoing testosterone suppression “were modest. The question of when it is fair to permit a transgender woman to compete in sport in line with her experienced gender identity is challenging.” (812)

164. Harper 2021: “[V]alues for strength, LBM [lean body mass], and muscle area in transwomen remain above those of cisgender women, even after 36 months of hormone therapy.” (1)

165. Hilton & Lundberg 2021: “evidence for loss of the male performance advantage, established by testosterone at puberty and translating in elite athletes to a 10–50% performance advantage, is lacking. . . . These data significantly

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

undermine the delivery of fairness and safety presumed by the criteria set out in transgender inclusion policies . . .” (211)

166. Hamilton et al. 2020, “Response to the United Nations Human Rights Council’s Report on Race and Gender Discrimination in Sport: An Expression of Concern and a Call to Prioritize Research”: “There is growing support for the idea that development influenced by high testosterone levels may result in retained anatomical and physiological advantages If a biologically male athlete self-identifies as a female, legitimately with a diagnosis of gender dysphoria or illegitimately to win medals, the athlete already possesses a physiological advantage that undermines fairness and safety. This is not equitable, nor consistent with the fundamental principles of the Olympic Charter.”

167. Hamilton et al. 2021, “Consensus Statement of the Fédération Internationale de Médecine du Sport” (International Federation of Sports Medicine, or FIMS), signed by more than 60 sports medicine experts from prestigious institutions around the world: The available studies “make it difficult to suggest that the athletic capabilities of transwomen individuals undergoing HRT or GAS are comparable to those of cisgender women.” The findings of Roberts et al. “question the required testosterone suppression time of 12 months for transwomen to be eligible to compete in women’s sport, as most advantages over ciswomen were not negated after 12 months of HRT.”

168. Outside the forum of peer-reviewed journals, respected voices in sport are reaching the same conclusion.

169. The **Women’s Sports Policy Working Group** identifies among its members and “supporters” many women Olympic medalists, former women’s tennis champion and LGBTQ activist Martina Navratilova, Professor Doriane Coleman, a former All-American women’s track competitor, transgender athletes Joanna Harper and Dr. Renee Richards, and many other leaders in women’s sports and civil rights. I have referenced other published work of Joanna Harper and Professor Coleman. In early 2021 the Women’s Sports Policy Working Group published a “Briefing Book” on the issue of transgender participation in women’s sports,¹³ in which they reviewed largely the same body of literature I have reviewed above, and analyzed the implications of that science for fairness and safety in women’s sports.

170. Among other things, the Women’s Sports Policy Working Group concluded:

¹³ <https://womenssportspolicy.org/wp-content/uploads/2021/02/Congressional-Briefing-WSPWG-Transgender-Women-Sports-2.27.21.pdf>

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

- “[T]he evidence is increasingly clear that hormones do not eliminate the legacy advantages associated with male physical development” (8) due to “the considerable size and strength advantages that remain even after hormone treatments or surgical procedures.” (17)
- “[T]here is convincing evidence that, depending on the task, skill, sport, or event, trans women maintain male sex-linked (legacy) advantages even after a year on standard gender-affirming hormone treatment.” (26, citing Roberts 2020.)
- “[S]everal peer-reviewed studies, including one based on data from the U.S. military, have confirmed that trans women retain their male sex-linked advantages even after a year on gender affirming hormones. . . . Because of these retained advantages, USA Powerlifting and World Rugby have recently concluded that it isn't possible fairly and safely to include trans women in women's competition.” (32)

171. As has been widely reported, in 2020, after an extensive scientific consultation process, the **World Rugby** organization issued its Transgender Guidelines, finding that it would not be consistent with fairness or safety to permit biological males to compete in World Rugby women's matches, no matter what hormonal or surgical procedures they might have undergone. Based on their review of the science, World Rugby concluded:

- “Current policies regulating the inclusion of transgender women in sport are based on the premise that reducing testosterone to levels found in biological females is sufficient to remove many of the biologically-based performance advantages described above. However, peer-reviewed evidence suggests that this is not the case.”
- “Longitudinal research studies on the effect of reducing testosterone to female levels for periods of 12 months or more do not support the contention that variables such as mass, lean mass and strength are altered meaningfully in comparison to the original male-female differences in these variables. The lowering of testosterone removes only a small proportion of the documented biological differences, with large, retained advantages in these physiological attributes, with the safety and performance implications described previously.”
- “. . . given the size of the biological differences prior to testosterone suppression, this comparatively small effect of testosterone reduction allows substantial and meaningful differences to remain. This has significant implications for the risk of injury”

- “. . . bone mass is typically maintained in transgender women over the course of at least 24 months of testosterone suppression, Height and other skeletal measurements such as bone length and hip width have also not been shown to change with testosterone suppression, and nor is there any plausible biological mechanism by which this might occur, and so sporting advantages due to skeletal differences between males and females appear unlikely to change with testosterone reduction.

172. In September 2021 the government-commissioned Sports Councils of the United Kingdom and its subsidiary parts (the five Sports Councils responsible for supporting and investing in sport across England, Wales, Scotland and Northern Ireland) issued a formal “Guidance for Transgender Inclusion in Domestic Sport” (UK Sport Councils 2021), following an extensive consultation process, and a commissioned “International Research Literature Review” prepared by the Carbmill Consulting group (UK Sport Literature Review 2021). The UK Sport Literature Review identified largely the same relevant literature that I review in this paper, characterizes that literature consistently with my own reading and description, and based on that science reaches conclusions similar to mine.

173. The UK Sport Literature Review 2021 concluded:

- “Sexual dimorphism in relation to sport is significant and the most important determinant of sporting capacity. The challenge to sporting bodies is most evident in the inclusion of transgender people in female sport.” “[The] evidence suggests that parity in physical performance in relation to gender-affected sport cannot be achieved for transgender people in female sport through testosterone suppression. Theoretical estimation in contact and collision sport indicate injury risk is likely to be increased for female competitors.” (10)
- “From the synthesis of current research, the understanding is that testosterone suppression for the mandated one year before competition will result in little or no change to the anatomical differences between the sexes, and a more complete reversal of some acute phase metabolic pathways such as haemoglobin levels although the impact on running performance appears limited, and a modest change in muscle mass and strength: The average of around 5% loss of muscle mass and strength will not reverse the average 40-50% difference in strength that typically exists between the two sexes.” (7)
- “These findings are at odds with the accepted intention of current policy in sport, in which twelve months of testosterone suppression is

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

expected to create equivalence between transgender women and females.” (7)

174. Taking into account the science detailed in the UK Sport Literature Review 2021, the UK Sports Councils have concluded:

- “[T]he latest research, evidence and studies made clear that there are retained differences in strength, stamina and physique between the average woman compared with the average transgender woman or non-binary person registered male at birth, with or without testosterone suppression.” (3)
- “Competitive fairness cannot be reconciled with self-identification into the female category in gender-affected sport.” (7)
- “As a result of what the review found, the Guidance concludes that the inclusion of transgender people into female sport cannot be balanced regarding transgender inclusion, fairness and safety in gender-affected sport where there is meaningful competition. This is due to retained differences in strength, stamina and physique between the average woman compared with the average transgender woman or non-binary person assigned male at birth, with or without testosterone suppression.” (6)
- “Based upon current evidence, testosterone suppression is unlikely to guarantee fairness between transgender women and natal females in gender-affected sports. . . . Transgender women are on average likely to retain physical advantage in terms of physique, stamina, and strength. Such physical differences will also impact safety parameters in sports which are combat, collision or contact in nature.” (7)

175. On January 15, 2022 the American Swimming Coaches Association (ASCA) issued a statement stating, “The American Swimming Coaches Association urges the NCAA and all governing bodies to work quickly to update their policies and rules to maintain fair competition in the women’s category of swimming. ASCA supports following all available science and evidenced-based research in setting the new policies, and we strongly advocate for more research to be conducted” and further stated “The current NCAA policy regarding when transgender females can compete in the women’s category can be unfair to cisgender females and needs to be reviewed and changed in a transparent manner.” (<https://swimswam.com/asca-issues-statement-calling-for-ncaa-to-review-transgender-rules/>; Accessed January 16, 2022.)

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

176. On January 19, 2022, the NCAA Board of Governors approved a change to the policy on transgender inclusion in sport and stated that “...the updated NCAA policy calls for transgender participation for each sport to be determined by the policy for the national governing body of that sport, subject to ongoing review and recommendation by the NCAA Committee on Competitive Safeguards and Medical Aspects of Sports to the Board of Governors. If there is no N[atational]G[overning]B[ody] policy for a sport, that sport's international federation policy would be followed. If there is no international federation policy, previously established IOC policy criteria would be followed”

(<https://www.ncaa.org/news/2022/1/19/media-center-board-of-governors-updates-transgender-participation-policy.aspx>; Accessed January 20, 2022.)

177. On February 1, 2022, because “...a competitive difference in the male and female categories and the disadvantages this presents in elite head-to-head competition ... supported by statistical data that shows that the top-ranked female in 2021, on average, would be ranked 536th across all short course yards (25 yards) male events in the country and 326th across all long course meters (50 meters) male events in the country, among USA Swimming members,” USA Swimming released its Athlete Inclusion, Competitive Equity and Eligibility Policy. The policy is intended to “provide a level-playing field for elite cisgender women, and to mitigate the advantages associated with male puberty and physiology.” (USA Swimming Releases Athlete Inclusion, Competitive Equity and Eligibility Policy, available at <https://www.usaswimming.org/news/2022/02/01/usa-swimming-releases-athlete-inclusion-competitive-equity-and-eligibility-policy>.) The policy states:

- For biologically male athletes seeking to compete in the female category in certain “elite” level events, the athlete has the burden of demonstrating to a panel of independent medical experts that:
 - “From a medical perspective, the prior physical development of the athlete as Male, as mitigated by any medical intervention, does not give the athlete a competitive advantage over the athlete’s cisgender Female competitors” and
 - There is a presumption that the athlete is not eligible unless the athlete “demonstrates that the concentration of testosterone in the athlete’s serum has been less than 5 nmol/L . . . continuously for a period of at least thirty-six (36) months before the date of the Application.” This presumption may be rebutted “if the Panel finds, in the unique circumstances of the case, that [the athlete’s prior physical development does not give the athlete a competitive advantage] notwithstanding the athlete’s serum testosterone results (e.g., the athlete has a medical condition

which limits bioavailability of the athlete's free testosterone)." (USA Swimming Athlete Inclusion Procedures at 43.)

Conclusions

The research and actual observed data show the following:

- At the level of (a) elite, (b) collegiate, (c) scholastic, and (d) recreational competition, men, adolescent boys, or male children, have an advantage over equally gifted, aged and trained women, adolescent girls, or female children in almost all athletic events;
- Biological male physiology is the basis for the performance advantage that men, adolescent boys, or male children have over women, adolescent girls, or female children in almost all athletic events; and
- The administration of androgen inhibitors and cross-sex hormones to men or adolescent boys after the onset of male puberty does not eliminate the performance advantage that men and adolescent boys have over women and adolescent girls in almost all athletic events. Likewise, there is no published scientific evidence that the administration of puberty blockers to males before puberty eliminates the pre-existing athletic advantage that prepubertal males have over prepubertal females in almost all athletic events.

For over a decade sports governing bodies (such as the IOC and NCAA) have wrestled with the question of transgender inclusion in female sports. The previous policies implemented by these sporting bodies had an underlying "premise that reducing testosterone to levels found in biological females is sufficient to remove many of the biologically-based performance advantages." (World Rugby 2020 at 13.) Disagreements centered around what the appropriate threshold for testosterone levels must be—whether the 10nmol/liter value adopted by the IOC in 2015, or the 5nmol/liter value adopted by the IAAF.

But the science that has become available within just the last few years contradicts that premise. Instead, as the UK Sports Councils, World Rugby, the FIMS Consensus Statement, and the Women's Sports Policy Working Group have all recognized the science is now sharply "at odds with the accepted intention of current policy in sport, in which twelve months of testosterone suppression is expected to create equivalence between transgender women and females" (UK Sports Literature Review 2021 at 7), and it is now "difficult to suggest that the athletic capabilities of transwomen individuals undergoing HRT or GAS are comparable to those of cisgender women." (Hamilton, FIMS Consensus Statement 2021.) It is important to note that while the 2021 "IOC Framework on Fairness,

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Inclusion, and Non-Discrimination on the Basis of Gender Identity and Sex Variations” calls for an “evidence-based approach,” that Framework does not actually reference *any* of the now extensive scientific evidence relating to the physiological differences between the sexes, and the inefficacy of hormonal intervention to eliminate male advantages relevant to most sports. Instead, the IOC calls on other sporting bodies to define criteria for transgender inclusion, while demanding that such criteria simultaneously ensure fairness, safety, and inclusion for all. The recently updated NCAA policy on transgender participation also relies on other sporting bodies to establish criteria for transgender inclusion while calling for fair competition and safety.

But what we currently know tells us that these policy goals—fairness, safety, and full transgender inclusion—are irreconcilable for many or most sports. Long human experience is now joined by large numbers of research papers that document that males outperform females in muscle strength, muscular endurance, aerobic and anaerobic power output, VO₂max, running speed, swimming speed, vertical jump height, reaction time, and most other measures of physical fitness and physical performance that are essential for athletic success. The male advantages have been observed in fitness testing in children as young as 3 years old, with the male advantages increasing immensely during puberty. To ignore what we know to be true about males’ athletic advantages over females, based on mere hope or speculation that cross sex hormone therapy (puberty blockers, androgen inhibitors, or cross-sex hormones) might neutralize that advantage, when the currently available evidence says it does not, is not science and is not “evidence-based” policy-making.

Because of the recent research and analysis in the general field of transgender athletics, many sports organizations have revised their policies or are in the process of doing so. As a result, there is not any universally recognized policy among sports organizations, and transgender inclusion policies are in a state of flux, likely because of the increasing awareness that the goals of fairness, safety, and full transgender inclusion are irreconcilable.

Sports have been separated by sex for the purposes of safety and fairness for a considerable number of years. The values of safety and fairness are endorsed by numerous sports bodies, including the NCAA and IOC. The existing evidence of durable physiological and performance differences based on biological sex provides a strong evidence-based rationale for keeping rules and policies for such sex-based separation in place (or implementing them as the case may be).

As set forth in detail in this report, there are physiological differences between males and females that result in males having a significant performance advantage over similarly gifted, aged, and trained females in nearly all athletic events before, during, and after puberty. There is not scientific evidence that any

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

amount or duration of cross sex hormone therapy (puberty blockers, androgen inhibitors, or cross-sex hormones) eliminates all physiological advantages that result in males performing better than females in nearly all athletic events. Males who have received such therapy retain sufficient male physiological traits that enhance athletic performance vis-à-vis similarly aged females and are thus, from a physiological perspective, more accurately categorized as male and not female.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Bibliography

- Bhargava, A. et al. *Considering Sex as a Biological Variable in Basic and Clinical Studies: An Endocrine Society Scientific Statement*. Endocr Rev. 42:219-258 (2021).
- Bohannon, R.W. et al., *Handgrip strength: a comparison of values obtained from the NHANES and NIH toolbox studies*. Am. J. Occ. Therapy 73(2) (March/April 2019).
- Catley, M. and G. Tomkinson, *Normative health-related fitness values for children: analysis of 85,437 test results on 9-17-year-old Australians since 1985*. Br. J. Sports Med. published online October 21, 2011. Bjsm.bmj.com. Additional versions of this article were published by BJSM in 2012 and 2013, including Br. J. Sports Med. 47:98-108 (2013).
- Chu, Y. et al., *Biomechanical comparison between elite female and male baseball pitchers*. J. App. Biomechanics 25:22-31 (2009).
- Coleman, D.L. and W. Shreve, *Comparing athletic performances: the best elite women to boys and men*. web.law.duke.edu/sites/default/files/centers/sportslaw/comparingathleticperformances.pdf. (Accessed 06/20/21)
- Coleman, D. L. et al., *Re-affirming the value of the sports exception to Title IX's general non-discrimination rule*. Duke J. of Gender and Law Policy 27(69):69-134 (2020).
- Davis SM, Kaar JL, Ringham BM, Hockett CW, Glueck DH, and Dabelea D. *Sex differences in infant body composition emerge in the first 5 months of life*. J Pediatr Endocrinol Metab 32: 1235-1239 (2019).
- De Miguel-Etayo, P. et al., *Physical fitness reference standards in European children: the IDEFICS study*. Int. J. Obes (Lond) 38(2):557-566 (2014).
- Dogan, B., *Multiple-choice reaction and visual perception in female and male elite athletes*. J. Sports Med. and Physical Fitness 49:91-96 (2009).
- Dykiert, D. and G. Der, *Sex differences in reaction time mean and intraindividual variability across the life span*. Developmental Psychology 48(5): 1262-76 (2012).
- Eiberg, S. et al, *Maximum oxygen uptake and objectively measured physical activity in Danish children 6-7 years of age: the Copenhagen school child intervention study*. Br J Sports Med 39:725-30 (2005).

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

- Fessler, D.M. et al. *Sexual dimorphism in foot length proportionate to stature*. Ann Hum Biol. 32:44-59 (2005).
- Fields J. et al., *Seasonal and Longitudinal Changes in Body Composition by Sport-Position in NCAA Division I Basketball Athletes*. Sports (Basel). 22:6 (2018).
- Gauthier. R. et al. The physical work capacity of Canadian children, aged 7 to 17 in 1983. A comparison with 1968. CAHPER Journal/Revue de l'ACSEPR 50:4–9 (1983).
- Gershoni, M. & Pietrokovski, S. *The landscape of sex-differential transcriptome and its consequent selection in human adults*. BMC BIOL 15: 7 (2017).
- Gooren, L., *The significance of testosterone for fair participation of the female sex in competitive sports*, 13 Asian J. of Andrology 653 (2011).
- Gooren, L., et al., *Transsexuals and competitive sports*. Eur. J. Endocrinol. 151:425-9 (2004).
- Haizlip, K.M. et al., *Sex-based differences in skeletal muscle kinetics and fiber-type composition*. PHYSIOLOGY (BETHESDA) 30: 30–39 (2015).
- Hamilton, B. et al, *Integrating transwomen and female athletes with differences of sex development (DSD) into elite competition: the FIMS 2021 consensus statement*. Sports Med 2021. doi: 10.1007/s40279-021-01451-8.
- Hamilton, B. et al., *Response to the United Nations Human Rights Council's report on race and gender discrimination in sport: an expression of concern and a call to prioritise research*. Sports Med 2020. doi: 10.1007/s40279-020-01380-4.
- Handelsman, D.J. et al., *Circulating testosterone as the hormonal basis of sex differences in athletic performance*. Endocrine Reviews 39(5):803-829 (Oct 2018).
- Handelsman, D.J., *Sex differences in athletic performance emerge coinciding with the onset of male puberty*, 87 Clinical Endocrinology 68 (2017).
- Handelsman, D.J., “Perspective,” at <https://www.healio.com/news/endocrinology/20201216/transgender-women-outpace-cisgender-women-in-athletic-tests-after-1-year-on-hormones> (last accessed September 29, 2021).
- Harper, J. et al., *How does hormone transition in transgender women change body composition, muscle strength and haemoglobin? Systematic review with a focus on the implications for sport participation*. Br J Sports Med 55(15):865-872 (2021).

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Harper, J., *Race time for transgender athletes*. J. Sporting Cultures & Identities 6:1 (2015).

Heydari R, et al. *Y chromosome is moving out of sex determination shadow*. Cell Biosci. 12:4. (2022).

Higerd, G.A. *Assessing the Potential Transgender Impact on Girl Champions in American High School Track and Field*. Doctoral Dissertation United States Sports Academy. (2020).
<https://www.proquest.com/openview/65d34c1e949899aa823beecad873afae/1?pq-origsite=gscholar&cbl=18750&diss=y>

Hilton, E. N. and T.R. Lundberg, *Transgender women in the female category of sport: perspectives on testosterone suppression and performance advantage*. Sports Medicine 51:199-214 (2021).

Hubal, M., H. Gordish-Dressman, P. Thompson, et al., *Variability in muscle size and strength gain after unilateral resistance training*, Med & Sci in Sports & Exercise 96:4 (2005).

Jain, A. et al., *A comparative study of visual and auditory reaction times on the basis of gender and physical activity levels of medical first year students*. Int J App & Basic Med Res 5:2(124-27) (May-Aug 2015).

Klaver M, et al.. *Early Hormonal Treatment Affects Body Composition and Body Shape in Young Transgender Adolescents*. J Sex Med 15: 251-260 (2018).

Knechtle, B., P. T. Nikolaidis et al., *World single age records in running from 5 km to marathon*, Frontiers in Psych 9(1) (2013).

Knox, T., L.C. Anderson, et al., *Transwomen in elite sport: scientific & ethical considerations*, 45 J. Med Ethics 395 (2019).

Lapauw, B. et al., *Body composition, volumetric and areal bone parameters in male-to-female transsexual persons*. Bone 43:1016-21 (2008).

Latorre-Roman, P. et al., *Reaction times of preschool children on the ruler drop test: a cross-sectional study with reference values*. Perceptual & Motor Skills 125(5):866-78 (2018).

Lepers, R., B. Knechtle et al., *Trends in triathlon performance: effects of sex & age*, 43 Sports Med 851 (2013).

Lesinski, M., A. Schmelcher, et al., *Maturation-, age-, and sex-specific anthropometric and physical fitness percentiles of German elite young athletes*. PLoS One. 15(8):e0237423 (2020).

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

- Levine, B., M. Joyner et al., *The role of testosterone in athletic performance*. Available at https://web.law.duke.edu/sites/default/files/centers/sportslaw/Experts_T_Statement_2019.pdf (January 2019).
- Leyk, D, W. Gorges et al., *Hand-grip strength of young men, women and highly trained female athletes*, Eur J Appl Physiol. 2007 Mar; 99(4):415-21 (2007).
- Lombardo, M. and R. Deaner, *On the evolution of the sex differences in throwing: throwing as a male adaptation in humans*, Quarterly Rev of Biology 93(2):91-119 (2018).
- Malina R.M., Bouchards, C., Bar-Or, O. Growth, Maturation, and Physical Activity (2nd edition). Published by Human Kinetics. 2004.
- McManus, A. and N. Armstrong, *Physiology of elite young female athletes*. J Med & Sport Sci 56:23-46 (2011).
- Millard-Stafford, M. et al., *Nature versus nurture: have performance gaps between men and women reached an asymptote?* Int'l J. Sports Physiol. & Performance 13:530-35 (2018).
- Miller, V.M. *Why are sex and gender important to basic physiology and translational and individualized medicine?* Am J Physiol Heart Circ Physiol 306(6): H781-788, (2014).
- Mormile, M. et al., *The role of gender in neuropsychological assessment in healthy adolescents*. J Sports Rehab 27:16-21 (2018).
- Morris, J. et al., *Sexual dimorphism in human arm power and force: implications for sexual selection on fighting ability*. J Exp Bio 223 (2020).
- National Collegiate Athletic Association, *Inclusion of transgender student-athletes*. https://ncaaorg.s3.amazonaws.com/inclusion/lgbtq/INC_TransgenderHandbook.pdf (August 2011).
- Neder, J.A. et al., *Reference values for concentric knee isokinetic strength and power in nonathletic men and women from 20 to 80 years old*. J. Orth. & Sports Phys. Therapy 29(2):116-126 (1999).
- Pate, R. and A. Kriska, *Physiological basis of the sex difference in cardiorespiratory endurance*. Sports Med 1:87-98 (1984).

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

- Pocek, S. et al., *Anthropometric Characteristics and Vertical Jump Abilities by Player Position and Performance Level of Junior Female Volleyball Players*. Int J Environ Res Public Health. 18: 8377-8386 (2021).
- Ramírez-Vélez, R. et al., *Vertical Jump and Leg Power Normative Data for Colombian Schoolchildren Aged 9-17.9 Years: The FUPRECOL Study*. J Strength Cond Res. 31: 990-998 (2017).
- Roberts, S.A., J.M. Carswell. *Growth, growth potential, and influences on adult height in the transgender and gender-diverse population*. Andrology. 9:1679-1688 (2021).
- Roberts, T.A. et al., *Effect of gender affirming hormones on athletic performance in transwomen and transmen: implications for sporting organisations and legislators*. Br J Sports Med published online at [10.1136/bjsports-2020-102329](https://doi.org/10.1136/bjsports-2020-102329) (Dec. 7, 2020).
- Roser, M., Cameron Appel and Hannah Ritchie (2013) "Human Height". Published online at OurWorldInData.org. Retrieved from: <https://ourworldindata.org/human-height> [Online Resource]
- Ross, J.G. and G.G. Gilbert. *National Children and Youth Fitness Study*. J Physical Educ Rec Dance (JOPERD) 56: 45 – 50 (1985).
- Sakamoto, K. et al., *Comparison of kicking speed between female and male soccer players*. Procedia Eng 72:50-55 (2014).
- Santos, R. et al. *Physical fitness percentiles for Portuguese children and adolescents aged 10-18 years*. J Sports Sci. 32:1510-8. (2014).
- Sattler, T. et al., *Vertical jump performance of professional male and female volleyball players: effects of playing position and competition level*. J Strength & Cond Res 29(6):1486-93 (2015).
- Sax L. *How common is intersex? a response to Anne Fausto-Sterling*. J Sex Res. 39(3):174-8 (2002).
- Scharff, M. et al., *Change in grip strength in trans people and its association with lean body mass and bone density*. Endocrine Connections 8:1020-28 (2019).
- Senefeld, J.W., et al. *Divergence in timing and magnitude of testosterone levels between male and female youths*, JAMA 324(1):99-101 (2020).
- Shah, K. et al. *Do you know the sex of your cells?* Am J Physiol Cell Physiol. 306(1):C3-18 (2014).

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

- Silverman, I., *The secular trend for grip strength in Canada and the United States*, J Sports Sci 29(6):599-606 (2011).
- Spierer, D. et al., *Gender influence on response time to sensory stimuli*. J Strength & Cond Res 24:4(957-63) (2010).
- Staiano AE, Katzmarzyk PT. *Ethnic and sex differences in body fat and visceral and subcutaneous adiposity in children and adolescents*. Int J Obes (Lond). 36:1261-9. (2012).
- Tack, L.J.W. et al., *Proandrogenic and antiandrogenic progestins in transgender youth: differential effects on body composition and bone metabolism*. J. Clin. Endocrinol. Metab, 103(6):2147-56 (2018).
- Tambalis, K. et al., *Physical fitness normative values for 6-18-year-old Greek boys and girls, using the empirical distribution and the lambda, mu, and sigma statistical method*. Eur J Sports Sci 16:6(736-46) (2016).
- Taylor, M.J. et al., *Vertical jumping and leg power normative data for English school children aged 10-15 years*. J Sports Sci. 28:867-72. (2010).
- Taylor RW, Gold E, Manning P, and Goulding A. *Gender differences in body fat content are present well before puberty*. Int J Obes Relat Metab Disord 21: 1082-1084, 1997.
- Taylor RW, Grant AM, Williams SM, and Goulding A. *Sex differences in regional body fat distribution from pre- to postpuberty*. Obesity (Silver Spring) 18: 1410-1416, 2010.
- Thibault, V., M. Guillaume et al., *Women and men in sport performance: the gender gap has not evolved since 1983*. J Sports Science & Med 9:214-223 (2010).
- Thomas, J.R. and K. E. French, *Gender differences across age in motor performance: a meta-analysis*. Psych. Bull. 98(2):260-282 (1985).
- Tomkinson, G. et al., *European normative values for physical fitness in children and adolescents aged 9-17 years: results from 2,779,165 Eurofit performances representing 30 countries*. Br J Sports Med 52:1445-56 (2018).
- Tomkinson, G. et al., *International normative 20 m shuttle run values from 1,142,026 children and youth representing 50 countries*. Br J Sports Med. 51:1545-1554 (2017).

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Tønnessen, E., I. S. Svendsen et al., *Performance development in adolescent track & field athletes according to age, sex, and sport discipline*. PLoS ONE 10(6): e0129014 (2015).

Tønnessen, E. et al., *Reaction time aspects of elite sprinters in athletic world championships*. J Strength & Cond Res 27(4):885-92 (2013).

United Kingdom Sports Councils, *Guidance for transgender inclusion in domestic sport*. Available at [https://equalityinsport.org/docs/300921/Guidance for Transgender Inclusion in Domestic Sport 2021 - Summary of Background Documents.pdf](https://equalityinsport.org/docs/300921/Guidance%20for%20Transgender%20Inclusion%20in%20Domestic%20Sport%202021%20-%20Summary%20of%20Background%20Documents.pdf). September 2021.

United Kingdom Sports Councils, *International Research Literature Review*. Available at <https://equalityinsport.org/docs/300921/Transgender%20International%20Research%20Literature%20Review%202021.pdf>. September 2021.

USA Swimming Athlete Inclusion Procedures, last revision February 1, 2022, available at https://www.usaswimming.org/docs/default-source/governance/governance-lsc-website/rules_policies/usa-swimming-policy-19.pdf.

VanCaenegem, E. et al, *Preservation of volumetric bone density and geometry in trans women during cross-sex hormonal therapy: a prospective observational study*. Osteoporos Int 26:35-47 (2015).

Wiik, A., T. R Lundberg et al., *Muscle strength, size, and composition following 12 months of gender-affirming treatment in transgender individuals*. J. Clinical Endocrin. & Metab. 105(3):e805-813 (2020).

Women's Sports Policy Working Group, *Briefing book: a request to Congress and the administration to preserve girls' and women's sport and accommodate transgender athletes*. Available at womenssportspolicy.org. (2021).

World Rugby Transgender Guidelines. <https://www.world.rugby/the-game/player-welfare/guidelines/transgender> (2020).

World Rugby Transgender Women's Guidelines. <https://www.world.rugby/the-game/player-welfare/guidelines/transgender/women> (2020).

Wunderlich RE, Cavanagh PR. *Gender differences in adult foot shape: implications for shoe design*. Med Sci Sports Exerc. 33:605-1 (2001).

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Appendix 1 – Data Tables**Presidential Physical Fitness Results¹⁴****Curl-Ups (# in 1 minute)**

Age	Male		Female		Age	Male-Female % Difference	
	50th %ile	85th %ile	50th %ile	85th %ile		50th %ile	85th %ile
6	22	33	23	32	6	-4.3%	3.1%
7	28	36	25	34	7	12.0%	5.9%
8	31	40	29	38	8	6.9%	5.3%
9	32	41	30	39	9	6.7%	5.1%
10	35	45	30	40	10	16.7%	12.5%
11	37	47	32	42	11	15.6%	11.9%
12	40	50	35	45	12	14.3%	11.1%
13	42	53	37	46	13	13.5%	15.2%
14	45	56	37	47	14	21.6%	19.1%
15	45	57	36	48	15	25.0%	18.8%
16	45	56	35	45	16	28.6%	24.4%
17	44	55	34	44	17	29.4%	25.0%

¹⁴ This data is available from a variety of sources, including:
<https://gilmore.gvdsd.us/documents/Info/Forms/Teacher%20Forms/Presidentialchallengephysicalfitness.pdf>

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Shuttle Run (seconds)

Age	Male		Female		Age	Male-Female % Difference	
	50th %ile	85th %ile	50th %ile	85th %ile		50th %ile	85th %ile
6	13.3	12.1	13.8	12.4	6	3.6%	2.4%
7	12.8	11.5	13.2	12.1	7	3.0%	5.0%
8	12.2	11.1	12.9	11.8	8	5.4%	5.9%
9	11.9	10.9	12.5	11.1	9	4.8%	1.8%
10	11.5	10.3	12.1	10.8	10	5.0%	4.6%
11	11.1	10	11.5	10.5	11	3.5%	4.8%
12	10.6	9.8	11.3	10.4	12	6.2%	5.8%
13	10.2	9.5	11.1	10.2	13	8.1%	6.9%
14	9.9	9.1	11.2	10.1	14	11.6%	9.9%
15	9.7	9.0	11.0	10.0	15	11.8%	10.0%
16	9.4	8.7	10.9	10.1	16	13.8%	13.9%
17	9.4	8.7	11.0	10.0	17	14.5%	13.0%

1 mile run (seconds)

Age	Male		Female		Age	Male-Female % Difference	
	50th %ile	85th %ile	50th %ile	85th %ile		50th %ile	85th %ile
6	756	615	792	680	6	4.5%	9.6%
7	700	562	776	636	7	9.8%	11.6%
8	665	528	750	602	8	11.3%	12.3%
9	630	511	712	570	9	11.5%	10.4%
10	588	477	682	559	10	13.8%	14.7%
11	560	452	677	542	11	17.3%	16.6%
12	520	431	665	503	12	21.8%	14.3%
13	486	410	623	493	13	22.0%	16.8%
14	464	386	606	479	14	23.4%	19.4%
15	450	380	598	488	15	24.7%	22.1%
16	430	368	631	503	16	31.9%	26.8%
17	424	366	622	495	17	31.8%	26.1%

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Pull Ups (# completed)

Age	Male		Female		Age	Male-Female % Difference	
	50th %ile	85th %ile	50th %ile	85th %ile		50th %ile	85th %ile
6	1	2	1	2	6	0.0%	0.0%
7	1	4	1	2	7	0.0%	100.0%
8	1	5	1	2	8	0.0%	150.0%
9	2	5	1	2	9	100.0%	150.0%
10	2	6	1	3	10	100.0%	100.0%
11	2	6	1	3	11	100.0%	100.0%
12	2	7	1	2	12	100.0%	250.0%
13	3	7	1	2	13	200.0%	250.0%
14	5	10	1	2	14	400.0%	400.0%
15	6	11	1	2	15	500.0%	450.0%
16	7	11	1	1	16	600.0%	1000.0%
17	8	13	1	1	17	700.0%	1200.0%

Data Compiled from Athletic.Net

2021 National 3000 m cross country race time in seconds

Rank	7-8 years old			9-10 years old			11-12 year old		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	691.8	728.4	Difference	607.7	659.8	Difference	608.1	632.6	Difference
2	722.5	739.0	#1 boy vs #	619.6	674.0	#1 boy vs #	608.7	639.8	#1 boy vs #
3	740.5	783.0	1 girl	620.1	674.7	1 girl	611.3	664.1	1 girl
4	759.3	783.5	5.0%	643.2	683.7	7.9%	618.6	664.4	3.9%
5	759.6	792.8		646.8	685.0		619.7	671.6	
6	760.0	824.1		648.0	686.4		631.2	672.1	
7	772.0	825.7	Average	648.8	687.0	Average	631.7	672.3	Average
8	773.0	832.3	difference	658.0	691.0	difference	634.9	678.4	difference
9	780.7	834.3	boys vs girls	659.5	692.2	boys vs girls	635.0	679.3	boys vs girls
10	735.1	844.4	6.2%	663.9	663.3	5.6%	635.1	679.4	6.3%

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

2021 National 3000 m cross country race time in seconds

Rank	5 th grade			6 th grade			7 th grade		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	625.5	667.0	Difference	545.3	582.0	Difference	534.0	560.7	Difference
2	648.8	685.0	#1 boy vs #	553.2	584.3	#1 boy vs #	541.0	567.0	#1 boy vs #
3	653.5	712.9	1 girl	562.3	585.1	1 girl	542.6	581.8	1 girl
4	658.4	719.2	6.2%	562.9	599.8	6.3%	544.6	583.0	4.8%
5	675.3	725.2		571.5	612.9		546.0	595.0	
6	677.4	727.7		588.0	622.0		556.0	599.0	
7	677.6	734.0	Average	591.3	624.9	Average	556.0	604.3	Average
8	679.1	739.4	difference	593.0	626.0	difference	556.0	606.0	difference
9	686.4	739.4	boys vs girls	593.8	628.0	boys vs girls	558.6	606.8	boys vs girls
10	686.4	746.4	7.3%	594.1	645.6	5.8%	563.2	617.0	7.1%

2021 National 100 m Track race time in seconds

Rank	7-8 years old			9-10 years old			11-12 year old		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	13.06	14.24	Difference #1	10.87	12.10	Difference #1	11.37	12.08	Difference #1
2	13.54	14.41	boy vs # 1	10.91	12.24	boy vs # 1	11.61	12.43	boy vs # 1
3	13.73	14.44	girl	11.09	12.63	girl	11.73	12.51	girl
4	14.10	14.48	8.3%	11.25	12.70	10.2%	11.84	12.55	5.9%
5	14.19	14.49		11.27	12.75		11.89	12.57	
6	14.31	14.58		11.33	12.80		11.91	12.62	
7	14.34	14.69	Average	11.42	12.83	Average	11.94	12.65	Average
8	14.35	14.72	difference	11.43	12.84	difference	11.97	12.71	difference
9	14.41	14.77	boys vs girls	11.44	12.88	boys vs girls	12.08	12.71	boys vs girls
10	14.43	14.86	3.6%	11.51	12.91	11.1%	12.12	12.75	5.7%

2021 National 200 m Track race time in seconds

Rank	7-8 years old			9-10 years old			11-12 year old		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	24.02	28.72	Difference #1	21.77	25.36	Difference #1	20.66	25.03	Difference #1
2	24.03	28.87	boy vs # 1	22.25	25.50	boy vs # 1	22.91	25.18	boy vs # 1
3	28.07	29.92	girl	22.48	25.55	girl	23.14	25.22	girl
4	28.44	29.95	16.4%	22.57	25.70	14.2%	23.69	25.49	17.5%
5	28.97	30.04		22.65	26.08		23.84	25.78	
6	29.26	30.09		22.77	26.22		24.23	25.89	
7	29.34	30.27	Average	23.11	26.79	Average	24.35	26.03	Average
8	29.38	30.34	difference	23.16	26.84	difference	24.58	26.07	difference
9	29.65	30.41	boys vs girls	23.28	26.91	boys vs girls	24.59	26.10	boys vs girls
10	29.78	30.54	6.1%	23.47	26.85	13.1%	24.61	26.13	7.9%

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

2021 National 400 m Track race time in seconds

Rank	7-8 years old			9-10 years old			11-12 year old		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	66.30	67.12	Difference #1	49.29	56.80	Difference #1	51.96	55.70	Difference #1
2	66.88	67.67	boy vs # 1	50.47	58.57	boy vs # 1	55.52	57.08	boy vs # 1
3	67.59	67.74	girl	52.28	60.65	girl	55.58	57.60	girl
4	68.16	68.26	1.2%	52.44	61.45	13.2%	55.59	57.79	6.7%
5	68.51	68.37		53.31	61.81		55.72	58.02	
6	69.13	71.02		53.65	62.03		55.84	58.25	
7	69.75	72.73	Average	53.78	62.32	Average	55.92	59.25	Average
8	69.80	73.25	difference	54.51	62.33	difference	57.12	59.27	difference
9	69.81	73.31	boys vs girls	55.84	62.34	boys vs girls	57.18	59.40	boys vs girls
10	70.32	73.48	2.4%	55.90	62.40	13.0%	57.22	59.49	4.2%

2021 National 800 m Track race time in seconds

Rank	7-8 years old			9-10 years old			11-12 year old		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	152.2	157.9	Difference #1	120.8	141.4	Difference #1	127.8	138.5	Difference #1
2	155.2	164.6	boy vs # 1	124.0	142.2	boy vs # 1	129.7	143.1	boy vs # 1
3	161.0	164.9	girl	125.1	148.8	girl	130.5	144.2	girl
4	161.1	165.9	3.6%	125.6	151.3	14.5%	133.2	144.2	7.7%
5	161.2	168.5		126.5	151.6		136.2	144.9	
6	161.6	169.9		136.5	152.5		136.5	145.0	
7	161.8	171.5	Average	137.1	153.1	Average	136.7	145.2	Average
8	162.2	173.1	difference	138.5	153.7	difference	136.7	145.6	difference
9	165.3	173.4	boys vs girls	139.5	153.8	boys vs girls	137.0	145.6	boys vs girls
10	166.9	174.7	4.5%	140.2	154.2	12.6%	137.9	145.8	6.9%

2021 National 1600 m Track race time in seconds

Rank	7-8 years old			9-10 years old			11-12 year old		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	372.4	397.6	Difference #1	307.4	319.3	Difference #1	297.3	313.8	Difference #1
2	378.3	400.9	boy vs # 1	313.7	322.2	boy vs # 1	298.4	317.1	boy vs # 1
3	378.4	405.6	girl	315.0	322.6	girl	307.0	319.9	girl
4	402.0	435.2	6.3%	318.2	337.5	3.7%	313.9	323.3	5.2%
5	406.4	445.0		318.4	345.2		319.2	325.3	
6	413.4	457.0		320.5	345.7		320.4	326.2	
7	457.4	466.0	Average	327.0	345.9	Average	321.1	327.0	Average
8	473.3	466.8	difference	330.3	347.1	difference	321.9	330.0	difference
9	498.3	492.3	boys vs girls	333.4	347.5	boys vs girls	325.5	331.1	boys vs girls
10	505.0	495.0	4.0%	347.0	355.6	4.7%	327.1	332.5	2.9%

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

2021 National 3000 m Track race time in seconds

Rank	7-8 years old			9-10 years old			11-12 year old		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	794.2	859.9	Difference #1	602.3	679.2	Difference #1	556.6	623.7	Difference #1
2	856.3		boy vs # 1	644.9	709.7	boy vs # 1	591.6	649.5	boy vs # 1
3			girl	646.6	714.2	girl	600.8	651.6	girl
4			7.6%	648.2	741.9	11.3%	607.1	654.9	10.8%
5				648.4	742.7		609.1	662.9	
6	No further data	No Further Data		652.8	756.6		611.5	664.1	
7			Average	658.9	760.2	Average	615.7	666.3	Average
8			difference	660.1	762.5	difference	617.3	666.8	difference
9			boys vs girls	662.7	780.2	boys vs girls	618.4	673.2	boys vs girls
10			NA%	671.6	792.3	12.7%	620.6	674.4	8.2%

2021 National Long Jump Distance (in inches)

Rank	7-8 years old			9-10 years old			11-12 year old		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	156.0	176.0	Difference #1	256.8	213.8	Difference #1	224.0	201.3	Difference #1
2	156.0	163.8	boy vs # 1	247.0	212.0	boy vs # 1	222.5	197.3	boy vs # 1
3	155.0	153.0	girl	241.0	210.8	girl	220.5	195.8	girl
4	154.3	152.0	-11.4%	236.3	208.8	20.1%	210.3	193.5	11.3%
5	154.0	149.5		231.5	207.0		210.0	193.3	
6	152.8	146.0		225.0	204.8		206.8	192.5	
7	151.5	144.5	Average	224.0	194.5	Average	206.0	192.3	Average
8	150.8	137.5	difference	224.0	192.5	difference	205.5	192.0	difference
9	150.5	137.0	boys vs girls	221.8	192.3	boys vs girls	205.0	191.3	boys vs girls
10		No Further Data	1.4%			13.2%			9.1%
	150.5			219.0	187.5		204.5	189.0	

2021 National High Jump Distance (in inches)

Rank	7-8 years old			9-10 years old			11-12 year old		
	Boys	Girls		Boys	Girls		Boys	Girls	
1	38.0	37.5	Difference #1	72.0	58.0	Difference #1	63.0	56.0	Difference #1
2	38.0	34.0	boy vs # 1	70.0	58.0	boy vs # 1	61.0	56.0	boy vs # 1
3	36.0	32.0	girl	65.8	57.0	girl	60.0	57.0	girl
4	36.0	32.0	1.3	62.0	56.0	24.1%	59.0	56.0	12.5%
5	35.8	32.0		62.0	56.0		59.0	56.0	
6	35.5			62.0	55.0		59.0	55.0	
7	34.0	No further Data	Average	61.0	54.0	Average	59.0	54.0	Average
8	32.0		difference	60.0	54.0	difference	58.0	54.0	difference
9	59.0		boys vs girls	59.0	No Further Data	boys vs girls	57.8	56.0	boys vs girls
10			21.6%			12.5%			6.9%
	56.0			56.0			57.8	56.0	

Appendix 2 – Scholarly Publications in Past 10 Years

Refereed Publications

1. Brown GA, Shaw BS, Shaw I. How much water is in a mouthful, and how many mouthfuls should I drink? A laboratory exercise to help students understand developing a hydration plan. *Adv Physiol Educ* 45: 589–593, 2021.
2. Schneider KM and Brown GA (as Faculty Mentor). What's at Stake: Is it a Vampire or a Virus? *International Journal of Undergraduate Research and Creative Activities*. 11, Article 4. 2019.
3. Christner C and Brown GA (as Faculty Mentor). Explaining the Vampire Legend through Disease. *UNK Undergraduate Research Journal*. 23(1), 2019. (*This is an on-campus publication.)
4. Schneekloth B and Brown GA. Comparison of Physical Activity during Zumba with a Human or Video Game Instructor. 11(4):1019-1030. *International Journal of Exercise Science*, 2018.
5. Bice MR, Hollman A, Bickford S, Bickford N, Ball JW, Wiedenman EM, Brown GA, Dinkel D, and Adkins M. Kinesiology in 360 Degrees. *International Journal of Kinesiology in Higher Education*, 1: 9-17, 2017
6. Shaw I, Shaw BS, Brown GA, and Shariat A. Review of the Role of Resistance Training and Musculoskeletal Injury Prevention and Rehabilitation. *Gavin Journal of Orthopedic Research and Therapy*. 1: 5-9, 2016
7. Kahle A, Brown GA, Shaw I, & Shaw BS. Mechanical and Physiological Analysis of Minimalist versus Traditionally Shod Running. *J Sports Med Phys Fitness*. 56(9):974-9, 2016
8. Bice MR, Carey J, Brown GA, Adkins M, and Ball JW. The Use of Mobile Applications to Enhance Learning of the Skeletal System in Introductory Anatomy & Physiology Students. *Int J Kines Higher Educ* 27(1) 16-22, 2016
9. Shaw BS, Shaw I, & Brown GA. Resistance Exercise is Medicine. *Int J Ther Rehab*. 22: 233-237, 2015.
10. Brown GA, Bice MR, Shaw BS, & Shaw I. Online Quizzes Promote Inconsistent Improvements on In-Class Test Performance in Introductory Anatomy & Physiology. *Adv. Physiol. Educ*. 39: 63-6, 2015
11. Brown GA, Heiserman K, Shaw BS, & Shaw I. Rectus abdominis and rectus femoris muscle activity while performing conventional unweighted and weighted seated abdominal trunk curls. *Medicina dello Sport*. 68: 9-18. 2015
12. Botha DM, Shaw BS, Shaw I & Brown GA. Role of hyperbaric oxygen therapy in the promotion of cardiopulmonary health and rehabilitation. *African Journal for*

Physical, Health Education, Recreation and Dance (AJPHERD). Supplement 2 (September), 20: 62-73, 2014

13. Abbey BA, Heelan KA, Brown, GA, & Bartee RT. Validity of HydraTrend™ Reagent Strips for the Assessment of Hydration Status. J Strength Cond Res. 28: 2634-9. 2014
14. Scheer KC, Siebrandt SM, Brown GA, Shaw BS, & Shaw I. Wii, Kinect, & Move. Heart Rate, Oxygen Consumption, Energy Expenditure, and Ventilation due to Different Physically Active Video Game Systems in College Students. International Journal of Exercise Science: 7: 22-32, 2014
15. Shaw BS, Shaw I, & Brown GA. Effect of concurrent aerobic and resistive breathing training on respiratory muscle length and spirometry in asthmatics. African Journal for Physical, Health Education, Recreation and Dance (AJPHERD). Supplement 1 (November), 170-183, 2013
16. Adkins M, Brown GA, Heelan K, Ansorge C, Shaw BS & Shaw I. Can dance exergaming contribute to improving physical activity levels in elementary school children? African Journal for Physical, Health Education, Recreation and Dance (AJPHERD). 19: 576-585, 2013
17. Jarvi MB, Brown GA, Shaw BS & Shaw I. Measurements of Heart Rate and Accelerometry to Determine the Physical Activity Level in Boys Playing Paintball. International Journal of Exercise Science: 6: 199-207, 2013
18. Brown GA, Krueger RD, Cook CM, Heelan KA, Shaw BS & Shaw I. A prediction equation for the estimation of cardiorespiratory fitness using an elliptical motion trainer. West Indian Medical Journal. 61: 114-117, 2013.
19. Shaw BS, Shaw I, & Brown GA. Body composition variation following diaphragmatic breathing. African Journal for Physical, Health Education, Recreation and Dance (AJPHERD). 18: 787-794, 2012.

Refereed Presentations

1. Brown GA. Transwomen competing in women's sports: What we know, and what we don't. American Physiological Society New Trends in Sex and Gender Medicine conference. Held virtually due to Covid-19 pandemic. October 19 - 22, 2021, 2021.
2. Shaw BS, Boshoff VE, Coetzee S, Brown GA, Shaw I. A Home-based Resistance Training Intervention Strategy To Decrease Cardiovascular Disease Risk In Overweight Children Med Sci Sport Exerc. 53(5), 742. 68th Annual Meeting of the American College of Sports Medicine. Held virtually due to Covid-19 pandemic. June 1-5, 2021.
3. Shaw I, Cronje M, Brown GA, Shaw BS. Exercise Effects On Cognitive Function And Quality Of Life In Alzheimer's Patients In Long-term Care. Med

- Sci Sport Exerc. 53(5), 743. 68th Annual Meeting of the American College of Sports Medicine. Held virtually due to Covid-19 pandemic. June 1-5, 2021.
4. Brown GA, Escalera M, Oleena A, Turek T, Shaw I, Shaw BS. Relationships between Body Composition, Abdominal Muscle Strength, and Well Defined Abdominal Muscles. Med Sci Sport Exerc. 53(5), 197. 68th Annual Meeting of the American College of Sports Medicine. Held virtually due to Covid-19 pandemic. June 1-5, 2021.
 5. Brown GA, Jackson B, Szekely B, Schramm T, Shaw BS, Shaw I. A Pre-Workout Supplement Does Not Improve 400 M Sprint Running or Bicycle Wingate Test Performance in Recreationally Trained Individuals. Med Sci Sport Exerc. 50(5), 2932. 65th Annual Meeting of the American College of Sports Medicine. Minneapolis, MN. June 2018.
 6. Paulsen SM, Brown GA. Neither Coffee Nor A Stimulant Containing “Pre-workout” Drink Alter Cardiovascular Drift During Walking In Young Men. Med Sci Sport Exerc. 50(5), 2409. 65th Annual Meeting of the American College of Sports Medicine. Minneapolis, MN. June 2018.
 7. Adkins M, Bice M, Bickford N, Brown GA. Farm to Fresh! A Multidisciplinary Approach to Teaching Health and Physical Activity. 2018 spring SHAPE America central district conference. Sioux Falls, SD. January 2018.
 8. Shaw I, Kinsey JE, Richards R, Shaw BS, and Brown GA. Effect Of Resistance Training During Nebulization In Adults With Cystic Fibrosis. International Journal of Arts & Sciences’ (IJAS). International Conference for Physical, Life and Health Sciences which will be held at FH Wien University of Applied Sciences of WKW, at Währinger Gürtel 97, Vienna, Austria, from 25-29 June 2017.
 9. Bongers M, Abbey BM, Heelan K, Steele JE, Brown GA. Nutrition Education Improves Nutrition Knowledge, Not Dietary Habits In Female Collegiate Distance Runners. Med Sci Sport Exerc. 49(5), 389. 64th Annual Meeting of the American College of Sports Medicine. Denver, CO. May 2017.
 10. Brown GA, Steele JE, Shaw I, Shaw BS. Using Elisa to Enhance the Biochemistry Laboratory Experience for Exercise Science Students. Med Sci Sport Exerc. 49(5), 1108. 64th Annual Meeting of the American College of Sports Medicine. Denver, CO. May 2017.
 11. Brown GA, Shaw BS, and Shaw I. Effects of a 6 Week Conditioning Program on Jumping, Sprinting, and Agility Performance In Youth. Med Sci Sport Exerc. 48(5), 3730. 63rd Annual Meeting of the American College of Sports Medicine. Boston, MA. June 2016.
 12. Shaw I, Shaw BS, Boshoff VE, Coetzee S, and Brown GA. Kinanthropometric Responses To Callisthenic Strength Training In Children. Med Sci Sport Exerc.

- 48(5), 3221. 63rd Annual Meeting of the American College of Sports Medicine. Boston, MA. June 2016.
13. Shaw BS, Shaw I, Gouveia M, McIntyre S, and Brown GA. Kinanthropometric Responses To Moderate-intensity Resistance Training In Postmenopausal Women. *Med Sci Sport Exerc.* 48(5), 2127. 63rd Annual Meeting of the American College of Sports Medicine. Boston, MA. June 2016.
 14. Bice MR, Cary JD, Brown GA, Adkins M, and Ball JW. The use of mobile applications to enhance introductory anatomy & physiology student performance on topic specific in-class tests. National Association for Kinesiology in Higher Education National Conference. January 8, 2016.
 15. Shaw I, Shaw BS, Lawrence KE, Brown GA, and Shariat A. Concurrent Resistance and Aerobic Exercise Training Improves Hemodynamics in Normotensive Overweight and Obese Individuals. *Med Sci Sport Exerc.* 47(5), 559. 62nd Annual Meeting of the American College of Sports Medicine. San Diego, CA. May 2015.
 16. Shaw BS, Shaw I, McCrorie C, Turner S., Schnetler A, and Brown GA. Concurrent Resistance and Aerobic Training in the Prevention of Overweight and Obesity in Young Adults. *Med Sci Sport Exerc.* 47(5), 223. 62nd Annual Meeting of the American College of Sports Medicine. San Diego, CA. May 2015.
 17. Schneekloth B, Shaw I, Shaw BS, and Brown GA. Physical Activity Levels Using Kinect™ Zumba Fitness versus Zumba Fitness with a Human Instructor. *Med Sci Sport Exerc.* 46(5), 326. 61st Annual Meeting of the American College of Sports Medicine. Orlando, FL. June 2014.
 18. Shaw I, Lawrence KE, Shaw BS, and Brown GA. Callisthenic Exercise-related Changes in Body Composition in Overweight and Obese Adults. *Med Sci Sport Exerc.* 46(5), 394. 61st Annual Meeting of the American College of Sports Medicine. Orlando, FL June 2014.
 19. Shaw BS, Shaw I, Fourie M, Gildenhuis M, and Brown GA. Variances In The Body Composition Of Elderly Woman Following Progressive Mat Pilates. *Med Sci Sport Exerc.* 46(5), 558. 61st Annual Meeting of the American College of Sports Medicine. Orlando, FL June 2014.
 20. Brown GA, Shaw I, Shaw BS, and Bice M. Online Quizzes Enhance Introductory Anatomy & Physiology Performance on Subsequent Tests, But Not Examinations. *Med Sci Sport Exerc.* 46(5), 1655. 61st Annual Meeting of the American College of Sports Medicine. Orlando, FL June 2014.
 21. Kahle, A. and Brown, G.A. Electromyography in the Gastrocnemius and Tibialis Anterior, and Oxygen Consumption, Ventilation, and Heart Rate During Minimalist versus Traditionally Shod Running. 27th National Conference on Undergraduate Research (NCUR). La Crosse, Wisconsin USA. April 11-13, 2013

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

22. Shaw, I., Shaw, B.S., and Brown, G.A. Resistive Breathing Effects on Pulmonary Function, Aerobic Capacity and Medication Usage in Adult Asthmatics Med Sci Sports Exerc 45 (5). S1602 2013. 60th Annual Meeting of the American College of Sports Medicine, Indianapolis, IN USA, May 26-30 2013
23. Shaw, B.S. Gildenhuis, G.A., Fourie, M. Shaw I, and Brown, G.A. Function Changes In The Aged Following Pilates Exercise Training. Med Sci Sports Exerc 45 (5). S1566 60th Annual Meeting of the American College of Sports Medicine, Indianapolis, IN USA, May 26-30 2013
24. Brown, G.A., Abbey, B.M., Ray, M.W., Shaw B.S., & Shaw, I. Changes in Plasma Free Testosterone and Cortisol Concentrations During Plyometric Depth Jumps. Med Sci Sports Exerc 44 (5). S598, 2012. 59th Annual Meeting of the American College of Sports Medicine. May 29 - June 2, 2012; San Francisco, California
25. Shaw, I., Fourie, M., Gildenhuis, G.M., Shaw B.S., & Brown, G.A. Group Pilates Program and Muscular Strength and Endurance Among Elderly Woman. Med Sci Sports Exerc 44 (5). S1426. 59th Annual Meeting of the American College of Sports Medicine. May 29 - June 2, 2012; San Francisco, California
26. Shaw B.S., Shaw, I., & Brown, G.A. Concurrent Inspiratory-Expiratory and Aerobic Training Effects On Respiratory Muscle Strength In Asthmatics. Med Sci Sports Exerc 44 (5). S2163. 59th Annual Meeting of the American College of Sports Medicine. May 29 - June 2, 2012; San Francisco, California
27. Scheer, K., Siebrandt, S., Brown, G.A, Shaw B.S., & Shaw, I. Heart Rate, Oxygen Consumption, and Ventilation due to Different Physically Active Video Game Systems. Med Sci Sports Exerc 44 (5). S1763. 59th Annual Meeting of the American College of Sports Medicine. May 29 - June 2, 2012; San Francisco, California
28. Jarvi M.B., Shaw B.S., Shaw, I., & Brown, G.A. (2012) Paintball Is A Blast, But Is It Exercise? Heart Rate and Accelerometry In Boys Playing Paintball. Med Sci Sports Exerc 44 (5). S3503. 59th Annual Meeting of the American College of Sports Medicine. May 29 - June 2, 2012; San Francisco, California

Book Chapters

1. Shaw BS, Shaw I, Brown G.A. Importance of resistance training in the management of cardiovascular disease risk. In Cardiovascular Risk Factors. IntechOpen, 2021.

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

2. Brown, G.A. Chapters on Androstenedione and DHEA. In: Nutritional Supplements in Sport, Exercise and Health an A-Z Guide. edited by Linda M. Castell, Samantha J. Stear, Louise M. Burke. Routledge 2015.

Refereed Web Content

1. Brown GA. Looking back and moving forward. The importance of reflective assessment in physiology education. (January 13, 2022)
<https://blog.lifescitrc.org/pecop/2022/01/13/looking-back-and-moving-forward-the-importance-of-reflective-assessment-in-physiology-education/>
2. Brown GA. The Olympics, sex, and gender in the physiology classroom. Physiology Educators Community of Practice, managed by the Education group of the American Physiological Society (August 18, 2021)
<https://blog.lifescitrc.org/pecop/2021/08/18/the-olympics-sex-and-gender-in-the-physiology-classroom/>

A complete CV is available at

https://www.unk.edu/academics/hperls/bio_pages/current-vita-gab.pdf